

ABSTRACT

304

PHYSICS / SCIENCE EDUCATION (PE)

ASSESSING CONCEPTUAL UNDERSTANDING OF ENERGY IN ELECTROSTATIC FIELDS: A CONCEPTUAL FRAMEWORK- BASED STUDY OF CHINESE HIGH SCHOOL STUDENTS

Jingyi Pang^{1,3}, Shihong Ma^{3,4}, Linghui Tian¹, Shuo Shen⁵, Lei Bao^{2,*}

¹ Institute of Higher Education, Fudan University, Shanghai 200433, China.

² School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing 211189, China.

³ Shanghai Research Center for Physics Education and Teaching, Shanghai 200433, China.

⁴ Department of Physics, Fudan University, Shanghai 200433, China.

⁵ High School Affiliated to Fudan University, Shanghai 200433, China.

* Contact author: lei.bao@outlook.com

ABSTRACT

In physics education reform, conceptual understanding is foundational to developing students' core physics literacy. However, students often rely on “problem-solving templates” for specific scenarios rather than achieving deep conceptual mastery. This challenge is particularly acute in understanding energy in electrostatic fields, where abstract and complex conceptual chains frequently lead to persistent misconceptions. This study develops a comprehensive conceptual framework for electrostatic energy that integrates physical laws, situational features, reasoning processes, and task objectives. Using this framework, we designed and validated a multiple-choice assessment tool to investigate students' understanding through knowledge integration and context dependency. Employing a mixed-methods approach with Chinese high school students who completed the electrostatic field unit, we identified and characterized cognitive profiles across three proficiency levels: Novice, Intermediate, and Expert. Empirical results reveal that the framework effectively characterizes significant differences in knowledge structures and thinking patterns across proficiency levels. Notably, most students exhibit substantial knowledge fragmentation and rely on memorization-based problem-solving, which impedes their ability to transfer concepts to novel contexts. These findings provide precise diagnostic information for physics instruction and offer significant pedagogical implications for optimizing instructional design of energy concepts in electrostatic fields.

REFERENCE

Student knowledge integration in learning mechanical wave propagation, *Physical Review Physics Education Research*, 17, 020122, 2021, L. Xie, Q. Liu, L. Hui.

Concentration analysis: A quantitative assessment of student states, *American Journal of Physics*, 69, S45–S53, 2001, L. Bao, E.F. Redish.

Teaching and assessing knowledge integration in science, *Science*, 313, 1049–1050, 2006, M.C. Linn, H. Lee, R. Tinker, F. Husic, J.L. Chiu.

Assessment of knowledge integration in student learning of galvanic cell: An interdisciplinary approach connecting physics and chemistry, *Physical Review Physics Education Research*, 20, 020129, 2024, D. Ye, J. Qian, W. Xu.

KEYWORDS

Conceptual understanding, Electrostatic field energy, Knowledge integration, Assessment tool, Physics education

HIDDEN RISKS AND OPTIMIZATION SCHEMES FOR APPLYING “SYMMETRY” IN HIGH SCHOOL PHYSICS TEACHING IN CHINA

X. Cui^{1,*}, S. Ma¹

¹Department of Physics, Fudan University, Shanghai.

ABSTRACT

In contemporary high school physics teaching in China, “as known from symmetry” is frequently taught to students as a convenient problem-solving skill to help them get answers faster. However, this oversimplified application which omits rigorous physical proofs easily traps students in a pattern of knowing the results without understanding the underlying mechanisms. Consequently, students become accustomed to relying on empirical intuition to comprehend physics, which hinders the development of physics ideas and scientific thinking. By providing rigorous physical proofs for common applications of symmetry in high school physics, and demonstrating symmetry from the perspective of physical laws rather than geometric intuition, teachers can expose students to the underlying logic of physics. This approach effectively fulfills the requirements for scientific thinking outlined in the physics core competencies in senior high schools. Furthermore, introducing analysis cases of non-ideal conditions and utilizing contrastive teaching methods can not only reinforce the prerequisites for the correct application of symmetry, but also highlight the rigorous logic inherent in physics, thereby further enhancing students’ analytical capabilities.

Abstract Topic: Physics and Science Education

REFERENCE

Explanation of the Physics Core Competencies and Instructional Suggestions in Senior High Schools, *Global Education*, 48(09), 77-88, 2019, B. Liao, H. Li, X. Li.

Comparison of University Students’ Understanding of Graphs in Different Contexts, *Physical Review Special Topics – Physics Education Research*, 9(2), 2013, M. Planinic, L. Ivanjek, A. Susac, Z. Milin-Sipus.

* Corresponding author email: 24210190006@m.fudan.edu.cn

Analysis and Comparison of Students' Conceptual Understanding of Symmetry Arguments in Gauss's and Ampere's Laws, *Physical Review Physics Education Research*, 19(1), 2023, E. Campos, E. Hernandez, P. Barniol, G. Zavala.

Using Symmetry and Invariance to Solve Problems in Elementary Physics, *The Physics Teacher*, 57, 475-477, 2019, M. Frodyma, S. Rosas.

Promoting the Construction of Intelligent Knowledge with the Help of Various Methods of Cognitively Activating Instruction, *Frontiers in Education*, 7: 979430, 2023, R. Schumacher, E. Stern.

KEYWORDS

Symmetry, Physics teaching, Scientific thinking

PRACTICAL EXPLORATION OF HIGH SCHOOL PHYSICS “PROJECT-BASED RESEARCH” WITH PBL-PAD CLASSROOM TEACHING MODEL

Zhiwu Xu^{1,2}, Shihong Ma^{2,3*}

¹ Institute of Higher Education, Fudan University, Shanghai 200433, China.

² Shanghai Research Center for Physics Education and Teaching, Shanghai 200433, China.

³ Department of Physics, Fudan University, Shanghai 200433, China.

ABSTRACT

The main content of this research is “Project-based research” to address the problems that exist universally in traditional experimental physics education. We created an innovative “PBL-PAD classroom” teaching model by combining practicability of project learning and the structural advantage of PAD (Presentation-Assimilation-Discussion) teaching model. This model has been explored through four stages of education.

This study adopts an experimental design. First, we chose M and N classes from F high school in Shanghai to conduct basic educational practice of PBL-PAD classroom research, and conducted a questionnaire survey. We used a five-point likert scale adapted from the ILS questionnaire to evaluate the learning style of students. Based on the classification of core physics literacy outlined in the *General High Schools Physics Curriculum Standards*, we used an evaluation scale to examine changes in core competence. The change of students’ physical and objective grades can be reflected through the test questions based on adaptation of relevant questions from the real questions of provincial and municipal examination or college entrance examination.

The results of the study showed that experimental and control groups had a clear difference in objective physical performance after receiving the PBL-PAD classroom teaching model. There is a clear hierarchical phenomenon in the total scores of class X’s and class Y’s physical core abilities, and there is no obvious difference between physical concept, scientific thinking, scientific attitude and responsibility. However, significant differences have been found at the level of scientific inquiry. In terms of learning style, class X do not differ significantly in the number of students showing an active and reflective learning style or a sensing and intuitive learning style. However, there are clear differences in the distributions of visual versus verbal learners and

* Corresponding author email: shma@fudan.edu.cn

sequential versus global learners, with the largest proportion of them being actively perceived.

REFERENCE STYLE

The use of project based learning to improve students' concept understanding on elasticity and Hooke law, *Asian Journal of Science Education*, 4, 1-8, 2022, SAFRIANA S, NURLINA N, GINTING F W, et al.

Project based learning and academic procrastination of students in learning physics, *International Journal of Instruction*, 13, 489-508, 2020, SANTYASA I W, RAPI N K, SARA I W W.

Effect of robotics-enhanced project-based learning approach on students' conceptual understanding and motivation in linear motion in physics in selected Rwandan Secondary School, *Education and information technologies*, 29, 12435-12456, 2024, GASANA J C, NKUNDABAKURA P, NSENGIMANA T, et al.

Learning and teaching styles in engineering education, *Engineering Education*, 78, 674, 1988, FELDER R M, SILVERMAN L K.

Reaching the second tier—learning and teaching styles in college science education, *Journal of college science teaching*, 22, 286-290, 1993, FELDER R M.

Applications, reliability, and validity of the index of learning styles, *Intl. J. Engr. Education*, 21, 103-112, 2005, FELDER R M, SPURLIN J E.

KEYWORDS

Project-based learning, PAD classroom, High school physics

THE DEVELOPMENT OF PHYSICS EDUCATION RESEARCH IN FUDAN UNIVERSITY

Shihong Ma^{1,2,*}, Lei Bao³

¹ Shanghai Research Center for Physics Education and Teaching, Shanghai 200433, China.

² Department of Physics, Fudan University, Shanghai 200433, China.

³ School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing 211189, China.

ABSTRACT

Physics education research is a discipline-based education research to explore practical problems in physics education, seek effective solutions, and improve the quality of physics education and teaching. The accumulated research results have had a profound impact on the construction of physics courses and teaching reforms. This presentation will present several typical examples to briefly introduce some of the research achievements in this field by Fudan University and its collaborating institutions in recent years. It also puts forward thoughts and suggestions for improving the current situation of physics education research in China, aiming to provide reference for promoting the development of domestic physics education research.

KEYWORDS

Physics education research, Research methods, Curriculum, Instructional resources

* Corresponding author email: shma@fudan.edu.cn

DIGITAL INTELLIGENCE DRIVING CLASSROOM REVOLUTION IN PHYSICS EDUCATION — A CASE STUDY OF THE BLENDED "SEMICONDUCTOR PHYSICS" COURSE

Fang Lu^{1,*}

¹ Department of Physics, Fudan University, Shanghai 200433, China.

ABSTRACT

In the era of digital intelligence, higher education is undergoing a profound shift from "teaching-centered" to "learning-centered" paradigms. Taking Fudan University's "Semiconductor Physics" course as an example, this report elaborates on how digital and intelligent technologies can be leveraged to establish a new teaching model. The course adopts a blended teaching model. In the online phase, we carry out problem-oriented learning, mind map construction, and online testing, to supplement students' knowledge gaps. Meanwhile, offline classrooms are transformed into student-led seminars, focusing on representative reports, critical discussions, in-class tests, and peer evaluations. The course assessment emphasizes process-oriented evaluation. Practice has shown that this hybrid design effectively enhances students' self-learning ability, higher-order thinking, and physics professional competence.

KEYWORDS

Digital Intelligence Driving, Physics Education, Blended Course, Semiconductor Physics

* Corresponding author email: fanglu@fudan.edu.cn

Opportunity or Challenge? Exploring Strategies for the Symbiotic Integration of GAI and Science Education

Zengze. Liu*, Leilei Pan, Xianglong Shan, Rutong Li

School of Physics and Technology, Soochow University, Suzhou, China

ABSTRACT

The rise of Generative Artificial Intelligence (GAI) presents both opportunities and challenges for a paradigm shift in science education. This study systematically examines the dual effects of integrating GAI, analyzes its enabling pathways and value-added potential, and critically evaluates the technical, ethical, and practical constraints. The study finds that GAI drives innovation through three pathways: in experimental design, by generating simulated environments that reduce cost risks and expand the boundaries of inquiry; in personalized instruction, by constructing adaptive learning pathways based on learning analytics to achieve data-driven differentiated instruction; and in promoting deep learning, by creating complex open-ended problem scenarios that guide students toward critical thinking and inquiry-based learning. This enhances the value of science education, facilitating interdisciplinary integration, personalized practice, and empowerment throughout the entire inquiry process.

However, the demands of science education for accuracy, rigor, and verifiability conflict with the limitations of GAI, such as “hallucinations,” uncertainty, and insufficient interpretability. Based on the current landscape, the study identifies practical challenges: insufficient technological applicability, particularly in high-precision scientific contexts; high implementation costs for teachers, including cognitive load, time investment, and lack of support; and data security and ethical issues involving privacy, algorithmic bias, and accountability. Therefore, there is an urgent need to develop a collaborative framework: establishing science-oriented quality standards and evaluation systems for GAI content; developing teacher professional development models centered on “human-machine collaboration” to strengthen digital literacy and technology integration capabilities; and formulating ethical governance and application guidelines to ensure technological reliability, fairness, and educational value. Through multidimensional advancement, we aim to build a new ecosystem for science education that is intelligent, trustworthy, and learner-centered.

* Corresponding author email: zengzeliu@suda.edu.cn

REFERENCE STYLE

- Enabling general chemistry students to take part in experimental design activities, *Chemistry Education Research and Practice*, 24(4): 1229-1242,2023,Scoggin J, Smith K C.
- Why Do Fishes Die? Lab Assignment Created Using ChatGPT, *Journal of Chemical Education*,101(8): 3171-3178,2024,Trčková K, Maršálek R, Žáček M, et al.
- An analysis of ai-generated laboratory reports across the chemistry curriculum and student perceptions of chatgpt,*Journal of Chemical Education*,100(11): 4351-4359,2023, West J K, Franz J L, Hein S M, et al.
- From data to noise to data for mixing physics across temperatures with generative artificial intelligence, *Proceedings of the National Academy of Sciences of the United States of America*, 119(32): e2203656119,2022,Yihang W, Lukas H, Pratyush T.
- Clark T M, Fhaner M, Stoltzfus M, et al. Using ChatGPT to Support Lesson Planning for the Historical Experiments of Thomson, Millikan, and Rutherford[J]. *Journal of Chemical Education*, 2024, 101(5): 1992-1999.
- Effective learning with a personal AI tutor: A case study,*Education and Information Technologies*,1-16,2024,Baillifard A, Gabella M, Lavenex P B, et al.
- Generative AI for Customizable Learning Experiences,*Sustainability*,16(7): 3034, 2024, Pesovski I, Santos R, Henriques R, et al.
- ChatGPT improves creative problem-solving performance in university students: An experimental study,*Computers & Education*,215: 105031,2024,Urban M, Děchtěrenko F, Lukavský J, et al.
- Performance of ChatGPT on the test of understanding graphs in kinematics,*Physical Review Physics Education Research*,20(1): 010109,2024,Polverini G, Gregorcic B.
- Using ChatGPT for Method Development and Green Chemistry Education in Upper-Level Laboratory Courses, *Journal of Chemical Education*,101(8): 3224-3232,2024,Ruff E F, Franz J L, West J K.
- Student interaction with ChatGPT can promote complex critical thinking skills, *Learning and Instruction*, 95: 102011,2025,Suriano R, Plebe A, Acciai A, et al.
- Exploring the AI-Human Interface for Personalized Learning in a Chemical Context, *Journal of Chemical Education*, 101(11): 4916-4923,2024,Clark T M, Tafini N.
- The influences of ChatGPT on undergraduate students' demonstrated and perceived interdisciplinary learning,*Education and Information Technologies*,1-27, 2024,Zhong T, Zhu G, Hou C, et al.
- Artificial Intelligence and Chemistry Teaching: A Propaedeutic Analysis of Chatgpt in Chemical Concepts Defining Br,*Quimica Nova*,46(9): 915-923.2023, Leite B S.
- The impact of students' use of ChatGPT on their research skills: The mediating effects of autonomous motivation, engagement, and self-directed learning, *Education and Information Technologies*, 1-32,2024,Li Y, Sadiq G, Qambar G, et al.
- Effectiveness of AI-assisted game-based learning on science learning outcomes, intrinsic motivation, cognitive load, and learning behavior[J]. *Education and Information Technologies*,1-22,2024,Chen C H, Chang C L.
- Can ChatGPT Enhance Chemistry Laboratory Teaching? Using Prompt Engineering to Enable AI in Generating Laboratory Activities,*Journal of Chemical Education*, 101(5): 1858-1864.2024,Araújo J L, Saúde I.
- How ChatGPT transformed teachers: the role of basic psychological needs in enhancing digital competence,*Frontiers in Psychology*, 15: 1458551,2024, Zhou J, Shen L, Chen W.

- How should we change teaching and assessment in response to increasingly powerful generative Artificial Intelligence? Outcomes of the ChatGPT teacher survey, *Education and Information Technologies*, 1-37, 2024, Bower M, Torrington J, Lai J W M, et al.
- Examining science education in ChatGPT: An exploratory study of generative artificial intelligence, *Journal of Science Education and Technology*, 32(3): 444-452, 2023, Cooper G.
- Confronting imminent challenges in humane epistemic agency in science education: An interview with ChatGPT, *Science & Education*, 1-27, 2024, Oh P S, Lee G G.
- Visualization of AI Accuracy: A Novel Assignment for the Teaching of Critical Thinking and Science Writing, *Teaching of Psychology*, 52(3): 285-290, 2024, Cail J.

KEYWORDS

Generative Artificial Intelligence; Science Education; Personalized Learning; Human-Machine Collaboration; Educational Ecosystem

DeepSeek-Assisted Physics Instructional Design: An Empirical Study of Teaching Newton's Third Law in a High School Setting

Dazhen Tong^{1,*}, Huihui Wu², A.Y.M. Atiquil Islam^{3,4}, Junci Liu², Lei Jiang⁵ and Hongmei Ren²

¹School of Teacher Education, Anhui Normal University, China.

²School of Physics and Electronic Information, Anhui Normal University, China.

³Joint Education Institute of Zhejiang Normal University and University of Kansas, Zhejiang Normal University China.

⁴China-Africa Regional Center for Digital Education Cooperation, Zhejiang Normal University, China.

⁵Tianjin Academy of Educational Science, China.

ABSTRACT

Generative artificial intelligence (GenAI) has shown growing potential in supporting teacher education, yet little is known about how pre-service teachers collaborate with AI during instructional design (ID). This study examined the effects of DeepSeek, a Chinese large language model, on pre-service physics teachers' ID performance and cognitive processes. A quasi-experimental design was conducted with 220 participants completing a 45-minute ID on Newton's Third Law, either with or without DeepSeek assistance. Data from IDs, interaction logs, and interviews were analyzed through statistical tests, Epistemic Network Analysis (ENA), and thematic coding. Results indicated that the DeepSeek-assisted group outperformed the control group in analytical and structural aspects of ID, such as Teaching Content Analysis and Learner Analysis. ENA revealed that high-performing participants exhibited proactive "problem representation–AI interaction–monitoring" patterns while low-performing participants relied on reactive prompting. Interviews further confirmed that AI was helpful in contextual introductions and experiment design but also revealed issues such as outdated curriculum references and overly absolute expressions that lacked sensitivity to teaching contexts. The findings suggest that GenAI can serve as a cognitive scaffold when used strategically and highlight the need to integrate AI literacy and metacognitive training into teacher education.

REFERENCE STYLE

[1] Visualising relativity: Assessing high school students' understanding of complex physics concepts through AI-generated, *Physics Education*, 59, 025018, 2024, M.G. de Souza, M. Won, D. Treagust, A. Serrano.

* Corresponding author email: dzton@ahnu.edu.cn

- [2] Leveraging AI for rapid generation of physics simulations in education: Building your own virtual lab, *The Physics Teacher*, 63, 424, 2025, Y. Ben-Zion, R.E. Zarzecki, J. Glazer, N.D. Finkelstein.
- [3] ChatGPT for good? On opportunities and challenges of large language models for education, *Learning and Individual Differences*, 103, 102274, 2023, E. Kasneci, S. Krebs, C. Spohn, F. Dorgerloh, A.-S. Mestres, C. Schulz.
- [4] ChatGPT in the higher education: A systematic literature review and research challenges, *International Journal of Educational Research*, 127, 102411, 2024, M.I. Baig, E. Yadegaridehkordi.
- [5] Students-Generative AI interaction patterns and its impact on academic writing, *Journal of Computing in Higher Education*, 1, 1, 2025, J. Kim, S.S. Lee, R. Detrick, J. Wang, N. Li.
- [6] Differences in student-AI interaction process on a drawing task: Focusing on students' attitude towards AI and the level of drawing skills, *Australasian Journal of Educational Technology*, 40, 19, 2024, J. Kim, Y. Ham, S.S. Lee.
- [7] Exploring the role of human-AI collaboration in solving scientific problems, *Physical Review Physics Education Research*, 21, 010149, 2025, D. Tong, B. Jin, Y. Tao, H. Ren, A.Y.M.A. Islam, L. Bao.
- [8] Exploring instructional design in K-12 STEM education: a systematic literature review, *International Journal of STEM Education*, 11, 43, 2024, S. Halawa, T.C. Lin, Y.S. Hsu.
- [9] A proficiency model for pre-service physics teachers' pedagogical content knowledge (PCK). What constitutes high-level PCK?, *Journal of Research in Science Teaching*, 60, 136, 2023, D. Schiering, S. Sorge, M.M. Keller, K. Neumann.
- [10] Professional knowledge affects action-related skills: The development of preservice physics teachers' explaining skills during a field experience, *Journal of Research in Science Teaching*, 57, 1554, 2020, C. Kulgemeyer, A. Borowski, D. Buschhüter, P. Enkrott, M. Kempin, P. Reinhold, C. Vogelsang.
- [11] A Review of Misconception in Physics: The Diagnosis, Causes, and Remediation, *Journal of Turkish Science Education*, 19, 403, 2022, G. Resbiantoro, R. Setiani, Dwikoranto.
- [12] Evaluating item difficulty patterns for assessing student misconceptions in science across physics, chemistry, and biology concepts, *Heliyon*, 7, e08388, 2021, S. Soeharto, B. Csapó.
- [13] A history of instructional design and its impact on educational psychology, *Historical Foundations of Educational Psychology*, -, 183, 1987, W. Dick.
- [14] Trends and Issues in Instructional Design, *Pearson Education*, 2, 11, 2007, R.A. Reiser, J.V. Dempsey.
- [15] Those who understand: Knowledge growth in teaching, *Educational Researcher*, 15, 4, 1986, L.S. Shulman.
- [16] From "basic knowledge and skills" to three-dimensional goals to core literacy: Three stages of curriculum and teaching reform in China over 40 years of reform and opening up, *Curriculum, Teaching Material and Method*, 39, 40, 2019, W.S. Yu.

- [17] From three-dimensional curriculum objectives to core literacy in physics, *Physics Teaching*, 39, 2, 2017, Y.Y. Guo.
- [18] Analysis and implications of the current situation of middle school physics teaching design from the perspective of core literacy: Based on an analysis of Physics Teacher articles from 2016 to 2020, *Physics Teacher*, 42, 63, 2021, N. Li, J. Zhang, Z. Tang.
- [19] Content-specific pedagogical knowledge, practices, and beliefs underlying the design of physics lessons: A case study, *Physical Review Physics Education Research*, 15, 010125, 2019, S. Kapon, A. Merzel.
- [20] Can ChatGPT support prospective teachers in physics task development?, *Physical Review Physics Education Research*, 19, 020128, 2023, S.K.S.S. Küchemann, S. Steinert, N. Revenga, M. Schweinberger, Y. Dinc, K.E. Avila, J. Kuhn.
- [21] Grading assistance for a handwritten thermodynamics exam using artificial intelligence: An exploratory study, *Physical Review Physics Education Research*, 20, 020144, 2024, G. Kortemeyer, J. Nöhl, D. Onishchuk.
- [22] Exploring generative AI assisted feedback writing for students' written responses to a physics conceptual question with prompt engineering and few-shot learning, *Physical Review Physics Education Research*, 20, 010152, 2024, T. Wang, Z. Chen.
- [23] Toward AI grading of student problem solutions in introductory physics: A feasibility study, *Physical Review Physics Education Research*, 19, 020163, 2023, G. Kortemeyer.
- [24] Using ChatGPT for teaching physics, *The Physics Teacher*, 62, 536, 2024, K.E. Avila, S. Steinert, S. Ruzika, J. Kuhn, S. Küchemann.
- [25] The use of ChatGPT in teaching and learning: A systematic review through SWOT analysis approach, *Frontiers in Education*, 9, 1328769, 2024, D.T.T. Mai, C.V. Da, N.V. Hanh.
- [26] Integrating artificial intelligence-based methods into qualitative research in physics education research: A case for computational grounded theory, *Physical Review Physics Education Research*, 19, 020123, 2023, P. Tschisgale, P. Wulff, M. Kubsch.
- [27] Human-AI interaction research agenda: A user-centered perspective, *Data and Information Management*, 8, 100078, 2024, T. Jiang, Z. Sun, S. Fu, Y. Lv.
- [28] Students' prompt patterns and its effects in AI-assisted academic writing: Focusing on students' level of AI literacy, *Journal of Research on Technology in Education*, 1, 1, 2025, J. Kim, S. Yu, S.S. Lee, R. Detrick.
- [29] Exploring artificial intelligence in academic essay: higher education student's perspective, *International Journal of Educational Research Open*, 5, 100296, 2023, A.R. Malik, Y. Pratiwi, K. Andajani, I.W. Numertayasa, S. Suharti, A. Darwis.
- [30] Human-AI collaboration patterns in AI-assisted academic writing, *Studies in Higher Education*, 49, 847, 2024, A. Nguyen, Y. Hong, B. Dang, X. Huang.
- [31] The effects of over-reliance on AI dialogue systems on students' cognitive abilities: a systematic review, *Smart Learning Environments*, 11, 28, 2024, C. Zhai, S. Wibowo, L. Li.

- [32] Using ChatGPT for science learning: A study on pre-service teachers' lesson planning, *IEEE Transactions on Learning Technologies*, 17, 1643, 2024, G.G. Lee, X. Zhai.
- [33] The Systematic Design of Instruction, *Pearson*, 7, -, 2008, W. Dick, L. Carey, J.O. Carey.
- [34] Using thematic analysis in psychology, *Qualitative Research in Psychology*, 3, 77, 2006, V. Braun, V. Clarke.
- [35] A guideline of selecting and reporting intraclass correlation coefficients for reliability research, *Journal of Chiropractic Medicine*, 15, 155, 2016, T.K. Koo, M.Y. Li.
- [36] Statistical Power Analysis for the Behavioral Sciences, *Lawrence Erlbaum Associates*, 2, -, 1988, J. Cohen.
- [37] Eta squared and partial eta squared as measures of effect size in educational research, *Educational Research Review*, 6, 135, 2011, J.T.E. Richardson.
- [38] Designing child-centric AI learning environments: Insights from LLM-enhanced creative project-based learning, *arXiv*, -, 2403.16159, 2024, S. Zha, Y. Qiao, Q. Hu, Z. Li, J. Gong, Y. Xu.
- [39] On the creativity of large language models, *AI and Society*, 40, 3785, 2025, G. Franceschelli, M. Musolesi.
- [40] Cognitive offloading, *Trends in Cognitive Sciences*, 20, 676, 2016, E.F. Risko, S.J. Gilbert.
- [41] Strategic offloading of delayed intentions into the external environment, *Quarterly Journal of Experimental Psychology*, 68, 971, 2015, S.J. Gilbert.

KEYWORDS

Generative artificial intelligence, Instructional design, DeepSeek, Newton's Third Law, Pre-service teachers

THREE FORMS OF ENACTED PCK IN PHYSICS TEACHING: TRANSFORMATIONS OF REPRESENTATION, LOGICAL STRUCTURE, AND MEANING OF PHYSICS KNOWLEDGE

Yi Ding^{1,*}, Huasen Zhou², Lei Bao³

¹School of Teacher Education, Nanjing University of Information Science and Technology, China.

²Yuqi Junior High School, China.

³School of Biomedical Engineering, Southeast University, China.

ABSTRACT

Traditionally, descriptive assessments of teachers' pedagogical content knowledge (PCK) have focused primarily on "what teachers ought to do" and "what teachers could do", while paying limited attention to "what teachers actually do" in classroom practice. Guided by the Refined Consensus Model (RCM), this study shifts the analytical focus toward the dynamic processes of teaching, examining how physics teachers mobilize, integrate, and transform subject matter knowledge in real time to address specific instructional challenges and achieve instructional goals. Based on an empirical analysis of multi-source data from 19 physics teachers, three categories of enacted PCK (ePCK) were identified: (1) transformation of the representation of subject matter knowledge, involving the activation of static symbols into dynamic sensory experiences; (2) transformation of the logical structure of subject matter knowledge, involving the integration of fragmented information into coherent conceptual frameworks and the reconstruction of knowledge production processes; and (3) transformation of the meaning of subject matter knowledge, involving the restoration of condensed disciplinary knowledge to real-life contexts and the design of structured cognitive conflicts to facilitate conceptual change. By further considering the quality of classroom interactions and students' learning outcomes, these ePCK forms were categorized into low-, medium-, and high-quality levels. This study advances PCK research toward a more practice-oriented approach that foregrounds the dynamic nature of teaching and teachers' professional agency, while offering both reflective insights and practical implications for improving physics teaching and student learning.

REFERENCE

Complementary Approaches to Teaching Nature of Science: Integrating Student Inquiry, Historical Cases, and Contemporary Cases in Classroom Practice. *Science Education*, 98(3), 461 – 486, 2014, Allchin, D., Andersen, H. M., & Nielsen, K.

* Corresponding author email: dingyi9701@163.com

Learning progressions as models and tools for supporting classroom assessment. Research Conference 2021: Excellent Progress for Every Student: Proceedings and Program, 47 – 51, 2021, Alonzo, A.

Declarative and Dynamic Pedagogical Content Knowledge as Elicited Through Two Video-Based Interview Methods, *Journal of Research in Science Teaching*, 53(8), 1259–1286, 2016, A.C. Alonzo, J. Kim.

Pedagogical Content Knowledge as Reflected in Teacher–Student Interactions: Analysis of Two Video Cases, *Journal of Research in Science Teaching*, 49(10), 1211–1239, 2012, A.C. Alonzo, M. Kobarg, T. Seidel.

Pre-service Chemistry Teachers' Use of Pedagogical Transformation Competence to Develop Topic-Specific Pedagogical Content Knowledge for Planning to Teach Acid–Base Equilibrium, *Chemistry Education Research and Practice*, 23(1), 137–158, 2022, S. Aydin-Gunbatar, F.N. Akin.

Use of Narrative Interviewing in Everyday Information Behavior Research, *Library & Information Science Research*, 26(1), 15–28, 2004, J.A. Bates.

Assessment and Measurement of Pedagogical Content Knowledge, in *Examining Pedagogical Content Knowledge*, 147–161, 1999, J.A. Baxter, N.G. Lederman.

On the Classification and Framing of Educational Knowledge, in *Knowledge and Control*, 47–69, 1971, B. Bernstein.

Construing Polanyi's Tacit Knowing as Knowing by Acquaintance Rather than Knowing by Representation: Some Implications, *The Polanyi Society Periodical*, 29(2), 26–43, 2003, D. Cannon.

The Refined Consensus Model of Pedagogical Content Knowledge in Science Education, in *Repositioning Pedagogical Content Knowledge in Teachers' Knowledge for Teaching Science*, 77–94, 2019, J. Carlson et al.

Investigating Practising Science Teachers' pPCK and ePCK Development as a Result of Collaborative CoRe Design, in *Repositioning Pedagogical Content Knowledge in Teachers' Knowledge for Teaching Science*, 225–252, 2019, J. Carpendale, A. Hume.

A Grand Rubric for Measuring Science Teachers' Pedagogical Content Knowledge, in *Repositioning Pedagogical Content Knowledge in Teachers' Knowledge for Teaching Science*, 253–271, 2019, K.K.H. Chan, M. Rollnick, J. Gess-Newsome.

Constructing CoRe as a Methodological for Capturing Pedagogical Content Knowledge: A Case Study of Thailand Teachers Teaching Global Warming, *Procedia - Social and Behavioral Sciences*, 116, 421–425, 2014, B. Chordnork, C. Yuenyong.

Thematic Analysis, *The Journal of Positive Psychology*, 12(3), 297–298, 2017, V. Clarke, V. Braun.

Using Bridging Analogies and Anchoring Intuitions to Deal with Students' Preconceptions in Physics, *Journal of Research in Science Teaching*, 30(10), 1241–1257, 1993, J. Clement.

Relationships of Knowledge and Practice: Teacher Learning in Communities, *Review of Research in Education*, 24(1), 249–305, 1999, M. Cochran-Smith, S.L. Lytle.

The Child and the Curriculum, in *John Dewey: The Middle Works, 1899–1924*, 1902/1976, J. Dewey.

Science as Subject-Matter and as Method, in *John Dewey: The Middle Works, 1899–1924*, 69–79, 1910/1978, J. Dewey.

Analysis of Students' Conceptual Change in Learning Newton's Third Law with an Integrated Framework of Model Analysis and Knowledge Integration, *Physical Review Physics Education Research*, 20(2), 020141, 2024, Y. Ding, G. Zhu, Q. Bian, L. Bao.

How to Study Thinking in Everyday Life: Contrasting Think-Aloud Protocols With Descriptions and Explanations of Thinking, *Mind, Culture, and Activity*, 5(3), 178–186, 1998, K.A. Ericsson, H.A. Simon.

Professional Knowledge of Science Teachers, in *Second International Handbook of Science Education*, 435–448, 2012, H.E. Fischer, A. Borowski, O. Tepner.

Attribution Error and the Quest for Teacher Quality, *Educational Researcher*, 39(8), 591–598, 2010, M.M. Kennedy.

Supporting Pre-service Life Sciences Teachers' Development of Pedagogical Content Knowledge Using Pre-recorded Teaching Videos, *Journal of Education*, (99), 189–211, 2025, H.C. Khoza.

Connecting Pedagogical Content Knowledge to Teaching Practice: Investigating Physics Teachers' Enacted PCK, Personal PCK, and Engagement with Student Ideas in Classroom Discussions, 2020, J. Kim.

Sample Size and Saturation in PhD Studies Using Qualitative Interviews, 11(3), 2010, M. Mason.

Cumulative and Segmented Learning: Exploring the Role of Curriculum Structures in Knowledge-Building, *British Journal of Sociology of Education*, 30(1), 43–57, 2009, K. Maton.

Making Semantic Waves: A Key to Cumulative Knowledge-Building, *Linguistics and Education*, 24(1), 8–22, 2013, K. Maton.

Exploring the Enactment of Pedagogical Content Knowledge about Electrostatics Through Teacher Talk by Two Pre-Service Teachers, *Research in Science Education*, 2025, E.N. Mazibe.

A Comparison Between Reported and Enacted Pedagogical Content Knowledge (PCK) About Graphs of Motion, *Research in Science Education*, 50(3), 941–964, 2020, E.N. Mazibe, C. Coetzee, E. Gaigher.

The Role of Representation in Teaching and Learning Critical Thinking, *Educational Review*, 54(1), 57–67, 2002, J. McKendree et al.

Technological Pedagogical Content Knowledge: A Framework for Teacher Knowledge, *Teachers College Record*, 108(6), 1017–1054, 2006, P. Mishra, M.J. Koehler.

Pedagogical Transformations of Science Content Knowledge in Korean Elementary Classrooms, *International Journal of Science Education*, 35(9), 1590–1624, 2013, P.S. Oh, K.S. Kim.

Unpacking PCK Relationships and Development in Science Teachers: A Systematic Synthesis of Empirical Research, S. Park, K.K.H. Chan.

The PCK Map Approach to Capturing the Complexity of Enacted PCK (ePCK) and Pedagogical Reasoning in Science Teaching, in *Repositioning Pedagogical Content Knowledge in Teachers' Knowledge for Teaching Science*, 187–199, 2019, S. Park, J.K. Suh.

Redesigning a First Year Physics Lab Course on the Basis of the Procedural and Conceptual Knowledge in Science Model, *Physical Review Physics Education Research*, 20(1), 010117, 2024, C.F.J. Pols, P.J.J.M. Dekkers.

The Cambridge Handbook of the Learning Sciences, Cambridge University Press, 2005, R.K. Sawyer.

Examining Assumptions about the Need for Teachers to Transform Subject Matter into Pedagogical Forms Accessible to Students, Teachers and Teaching, 28(1), 1–11, 2022, T. Scheiner.

Understanding and Supporting Science Teachers Teaching Outside Their Expertise: Studies of Teachers and Those Who Support Them, 2021, H. Singh.

Development of Enacted Pedagogical Content Knowledge (ePCK) Among Pre-service Chemistry Teachers, Chemistry Education Research and Practice, 26(3), 761–779, 2025, C. Yildiz et al.

Design-Based STEM Activities in Teacher Education and Its Effect on Pre-service Science Teachers' Design Thinking Skills, Journal of Science Education and Technology, 34(4), 904–918, 2025, A.O. Yüksel.

KEYWORDS

EPCK, Transformation of subject matter knowledge, RCM, Physics teacher

COMPARATIVE ANALYSIS OF THE NEW EDITION GRADE 7 SCIENCE TEXTBOOKS: EVIDENCE FROM LLM- AND HUMAN-BASED EVALUATION

Shujing Wu¹

¹School of Biological Science and Medical Engineering, Southeast University, Nanjing 211189, China

ABSTRACT

Traditional textbook evaluation methods face growing challenges as revision cycles shorten. Large language models (LLMs), with their capacities for semantic understanding and logical reasoning, offer a promising alternative; however, their reliability for such tasks remains insufficiently validated. This study employed both human raters and an LLM (Gemini-3-pro) to quantitatively analyze STEM-related content in the latest science textbooks from three major Chinese publishers. Human evaluation reveals that all three editions are science-centered in their STEM content distribution, with technology and mathematics primarily serving supportive roles. Engineering content shows the most notable variation across editions. Three-discipline integration is the most common interdisciplinary pattern. Comparing human and LLM performance, Gemini-3-pro demonstrates reproducible stability across repeated trials and achieves high precision but low recall, reflecting a conservative coding tendency. The model performs well when textual cues are explicit, but tends to miss content requiring contextual interpretation, particularly in the technology and mathematics dimensions, due to its reliance on surface-level semantic matching. Based on these findings, we propose a human–AI collaborative framework in which LLMs handle structured extraction and preliminary annotation, while human raters focus on boundary-ambiguous segments, thereby improving both efficiency and reliability.

KEYWORDS

STEM education, LLM, Science textbooks, Textbook analysis

LARGE LANGUAGE MODELS IN PHYSICS EDUCATION: FACTORS SHAPING UNCERTAINTY IN EXPLAINING PHYSICS PROBLEMS

Z. Fang¹, L. Bao¹

¹School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing, Jiangsu 211189, China.

ABSTRACT

Large language models (LLMs) show considerable promise for educational question answering and learning support, yet their application in physics education remains constrained by output uncertainty. Physics problems involve strict conditional constraints, well-defined conceptual boundaries, and multi-step reasoning, making LLM-generated explanations particularly sensitive to subtle variations in external settings. Consequently, the same problem may elicit substantially different responses under different conditions, raising concerns about reliability and interpretability. This study investigates the factors shaping LLM uncertainty in explaining physics problems, focusing on prompt design, retrieval-augmented generation (RAG), and temperature settings. We systematically examine how these variables influence response uncertainty and use visualization methods to illustrate variation patterns across conditions. This research provides a basis for evaluating the quality of LLM applications in physics education, optimizing model deployment strategies, and improving intelligent instructional systems for discipline-specific learning contexts.

REFERENCE STYLE

Naive beliefs in 'sophisticated' subjects: Misconceptions about trajectories of objects, *Cognition*, 9, 117-123, 1981, A. Caramazza, M. McCloskey, B. Green.

Three types of conceptual change: Belief revision, mental model transformation, and categorical shift, *International handbook of research on conceptual change*, 61-82, 2008, M.T.H. Chi.

Synthesis of discipline-based education research in physics, *Physical Review Special Topics: Physics Education Research*, 10, 020119, 2014, J.L. Docktor, J.P. Mestre.

LM-polygraph: Uncertainty estimation for language models, arXiv preprint arXiv:2311.07383, 2023, A. Fadeeva, A. Rubashevskii, A. Shelmanov, S. Petrakov, I. Khan, E. Tutubalina, A. Panchenko.

Detecting hallucinations in large language models using semantic consistency, *Nature*, 630, 625-630, 2024, S. Farquhar, J. Kossen, L. Kuhn, Y. Gal.

A survey of confidence estimation and calibration in large language models, *Proceedings of the North American Chapter of the Association for Computational Linguistics*, 6577-6595, 2024, J. Geng, F. Cai, Y. Wang, H. Koepl, P. Nakov, I. Gurevych.

On calibration of modern neural networks, *Proceedings of the International Conference on Machine Learning*, 1321-1330, 2017, C. Guo, G. Pleiss, Y. Sun, K.Q. Weinberger.

More than misconceptions: Multiple perspectives on student knowledge and reasoning, *American Journal of Physics*, 64, 1316-1325, 1996, D. Hammer.

Aleatory and epistemic uncertainty in probability elicitation with an example from hazardous waste management, *Reliability Engineering and System Safety*, 54, 217-223, 1996, S.C. Hora.

Aleatoric and epistemic uncertainty in machine learning: An introduction to concepts and methods, *Machine Learning*, 110, 457-506, 2021, E. Hullermeier, W. Waegeman.

Language models (mostly) know what they know, arXiv preprint arXiv:2207.05221, 2022, S. Kadavath et al.

ChatGPT for good? On opportunities and challenges of large language models for education, *Learning and Individual Differences*, 103, 102274, 2023, E. Kasneci et al.

Naive theories of motion, *Mental Models*, 299-324, 1983, M. McCloskey.

Rethinking the role of demonstrations: What makes in-context learning work?, *Proceedings of the Conference on Empirical Methods in Natural Language Processing*, 11048-11064, 2022, S. Min et al.

Generative large language models for dialog-based tutoring, *Proceedings of the International Conference on Educational Data Mining*, 548-554, 2023, B. Nye, D. Mee, M.G. Core.

Self-consistency improves chain of thought reasoning in language models, *International Conference on Learning Representations*, 2023, X. Wang et al.

Can LLMs express their uncertainty? An empirical evaluation of confidence elicitation in large language models, International Conference on Learning Representations, 2024, M. Xiong et al.

Calibrate before use: Improving few-shot performance of language models, Proceedings of the International Conference on Machine Learning, 12697-12706, 2021, Z. Zhao, E. Wallace, S. Feng, D. Klein, S. Singh.

KEYWORDS

physics education, large language models, uncertainty analysis, prompt design, retrieval augmentation

Corresponding author email: zijingfang@seu.edu.cn

ASSESSMENT OF KNOWLEDGE INTEGRATION IN STUDENT LEARNING OF GALVANIC CELL: AN INTERDISCIPLINARY APPROACH CONNECTING PHYSICS AND CHEMISTRY

D. Ye¹, L. Bao¹

¹School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing, Jiangsu 211189, China.

ABSTRACT

The galvanic cell is a typical interdisciplinary topic, which is primarily taught in chemistry but its underpinning is closely related to physics. Student learning in galvanic cell has been extensively studied in chemistry education, which has revealed a large number of misconceptions that are difficult to change through traditional instruction. A source of the learning difficulties is that most students lack an explanatory framework to integrate the many complex phenomena and processes into a coherent knowledge system. As a result, students often rely on memorization of terms, laws, and equations in solving problems but without meaningful understanding in their reasoning. To address the deficit in learning, this study aims to help students develop an explanatory framework for supporting an integrated knowledge structure. Specifically, a conceptual framework model about galvanic cell is developed based on the central idea from physics, which is used as the core concept for mechanistic explanations of the related chemistry concepts. Guided by the conceptual framework, an instrument is developed to assess the knowledge integration in student learning of galvanic cell. A combination of qualitative and quantitative measures were used to assess the levels of knowledge integration in learning galvanic cell with a large number of Chinese high school and college students. The assessment results show that the conceptual framework model can effectively represent the knowledge structures of students at different levels of knowledge integration. The galvanic cell conceptual framework and assessment results can provide useful resources for teachers to develop instruction for promoting knowledge integration.

REFERENCE STYLE

Paper Title, Journal Name, Volume Number, Page Numbers, Year of Publication, Initial1. LastName1, Initial2. LastName2, Initial3. LastName3, Initial4. LastName4.
Example: High-Performance Lithium-Ion Batteries Using Nanostructured Silicon Anodes, *Advanced Energy Materials*, 12, 2103456–2103465, 2022, J.A. Smith, L.B. Chen, M.K. Patel, R.T. Johnson.

REFERENCES

1. Approaches to understanding and measuring interdisciplinary scientific research (IDR): A review of the literature, *Journal of informetrics*, 5, 14-26, 2011, C. S. Wagner et al.
2. Integrated knowledge content in an interdisciplinary field: identification, classification, and application, *Scientometrics*, 127, 6581-6614, 2022, S. Wang, J. Mao, Y. Cao, G. Li.

3. Interdisciplinarity in science: A tentative typology of disciplines and research areas, *Journal of the American Society for Information Science and technology*, 54, 1237-1249, 2003, F. Morillo, M. Bordons, I. Gómez.
4. Diversity and network coherence as indicators of interdisciplinarity: case studies in bionanoscience, *Scientometrics*, 82, 263-287, 2010, I. Rafols, M. Meyer.
5. Taxonomy of educational objectives: the classification of educational goals: handbook I: cognitive domain, New York, US: D. McKay, 1956, B. S. Bloom, M. D. Engelhart, E. J. Furst, W. H. Hill, D. R. Krathwohl.
6. A taxonomy for learning, teaching, and assessing: A revision of Bloom's taxonomy of educational objectives complete edition, JSTOR, 2005, J. Conklin.
7. Investigating Differences in Experts' Chemistry Knowledge Structures and Comparing Them to Those of General Chemistry Students, *Journal of Chemical Education*, 99, 2950-2963, 2022, O. Gulacar et al.
8. Students' representations of the atomic structure—the effect of some individual differences in particular task contexts, *Chemistry Education Research and Practice*, 17, 209-219, 2016, G. Papageorgiou, A. Markos, N. Zarkadis.
9. Exploring conceptual frameworks of models of atomic structures and periodic variations, chemical bonding, and molecular shape and polarity: a comparison of undergraduate general chemistry students with high and low levels of content knowledge, *Chemistry Education Research and Practice*, 14, 130-146, 2013, C.-Y. Wang, L. H. Barrow.
10. Model analysis: Representing and assessing the dynamics of student learning, *Physical Review Special Topics-Physics Education Research*, 2, 010103, 2006, L. Bao, E. F. Redish.
11. Procedural and conceptual knowledge of expert and novice students for the solving of a basic problem in chemistry, *International Journal of Science Education*, 21, 195-211, 1999, R. M. Heyworth.
12. How experts differ from novices in How People Learn, National Academy Press, Washington DC, 1999, J. Bransford, A. Brown, R. Cocking.
13. Conceptual framework assessment of knowledge integration in student learning of measurement uncertainty, *Physical Review Physics Education Research*, 19, 2023, C. Lu et al.
14. Knowledge integration in student learning of Newton's third law: Addressing the action-reaction language and the implied causality, *Physical Review Physics Education Research*, 17, 020116, 2021, L. Bao, J. C. Fritchman.
15. The value of vague ideas in the development of the periodic system of chemical elements, *Synthese*, 199, 10587-10614, 2021, T. Vogt.
16. Modeling chemical reactions with the Gaussian gun, *Journal of Chemical Education*, 96, 100-103, 2018, L. A. Elliott, E. Sippola, J. Watkins.
17. The trouble with chemical energy: Why understanding bond energies requires an interdisciplinary systems approach, *CBE—Life Sciences Education*, 12, 306-312, 2013, M. M. Cooper, M. W. Klymkowsky.
18. Assessment of student knowledge integration in learning work and mechanical energy, *Physical Review Physics Education Research*, 19, 010127, 2023, D. Tong et al.
19. Student knowledge integration in learning mechanical wave propagation, *Physical Review Physics Education Research*, 17, 020122, 2021, L. Xie et al.
20. Assessment of knowledge integration in student learning of simple electric circuits, *Physical Review Physics Education Research*, 18, 020102, 2020, Z. Liu, S. Pan, X. Zhang, L. Bao.

21. Knowledge integration in student learning of Newton's third law: Addressing the action-reaction language and the implied causality, *Physical Review Physics Education Research*, 17, 020116, 2011, L. Bao, J. C. Fritchman.
22. Assessment of knowledge integration in student learning of momentum, *Physical Review Physics Education Research*, 16, 010130, 2020, W. Xu et al.
23. Assessment of student understanding on light interference, *Physical Review Physics Education Research*, 15, 020134, 2013, R. Dai, J. C. Fritchman, Q. Liu, Y. Xiao, H. Yu, L. Bao.
24. Teaching towards knowledge integration in learning force and motion, *International Journal of Science Education*, 41, 2271-2295, 2019, Y. Nie et al.
25. Around the world in electrochemistry: a review of the electrochemistry curriculum in high schools, *Journal of Solid State Electrochemistry*, 28, 1361-1374, 2024, K. L. Turner et al.
26. Assessment of electrochemical concepts: a comparative study involving senior high-school students in Indonesia and Japan, *Research in Science & Technological Education*, 29, 169-188, 2011, S. Rahayu, D. F. Treagust, A. Chandrasegaran, M. Kita, S. Ibnu.
27. *Chemistry: Principles of Chemical Reactions*, Beijing: People's education press, 93-103, 2019, J. Wang, C. Zheng.
28. *Chemistry: Theoretical Edition*, TOKYO: TOKYO SHOSEKI CO., LTD, 112-119, 2023, O. Keiichiro, M. Motoyuki.
29. Conceptual difficulties experienced by senior high school students of electrochemistry: Electrochemical (galvanic) and electrolytic cells, *Journal of Research in Science Teaching*, 29, 1079-1099, 1992, P. J. Garnett, D. F. Treagust.
30. Development of a measurement instrument to assess students' proficiency levels regarding galvanic cells, *Chemistry Education Research and Practice*, 21, 655-667, 2020, H. Lu, Y. Jiang, H. Bi.
31. L. Dingrando, N. Hainen, C. Wistrom, and D. Zike, *Glencoe Chemistry: Matter & Change*. Houston, TX: McGraw Hill Education, 708-727, 2017, T. Buthelezi.
32. *Heinemann CHEMISTRY 2*, Melbourne, Victoria: Pearson Australia, 125-168, 2017, L. Derry et al.
33. Effect of cooperative learning strategies on students' understanding of concepts in electrochemistry, *International Journal of science and mathematics education*, 5, 349-373, 2007, B. Acar, L. Tarhan.
34. High school students' learning difficulties in electrochemistry: a mini review, *African Journal of Chemical Education*, 12, 202-237, 2022, M. T. Ali, A. R. Woldu, A. G. Yohannes.
35. Effectiveness of conceptual change-oriented teaching strategy to improve students' understanding of galvanic cells, *Journal of Chemical Education*, 83, 1719, 2006, A. R. Özkaya, M. Üce, H. Saricayir, M. Sahin.
36. Conceptual difficulties experienced by prospective teachers in electrochemistry: Half-cell potential, cell potential, and chemical and electrochemical equilibrium in galvanic cells, *Journal of Chemical Education*, 79, 735, 2002, A. R. Özkaya.
37. Determining of the prospective teachers' understandings of electrochemistry, *Procedia-Social and Behavioral Sciences*, 174, 2831-2838, 2015, A. Yilmaz, S. Bayrakçeken.
38. Students' misconceptions in electrochemistry regarding current flow in electrolyte solutions and the salt bridge, *Journal of chemical education*, 74, 819, 1997, M. J. Sanger, T. J. Greenbowe.
39. The development and implementation of a two-tier multiple choice test related to the concepts of oxidation-reduction reactions and galvanic cells, *Journal of Educational Sciences & Psychology*, 12 (74), 34-47, 2022, A. R. Şekerci, A. R. Erdem.

40. Conceptual difficulties experienced by senior high school students of electrochemistry: Electric circuits and oxidation-reduction equations, *Journal of Research in Science Teaching*, 29, 121-142, 1992, P. J. Garnett, D. F. Treagust.
41. Common student misconceptions in electrochemistry: Galvanic, electrolytic, and concentration cells, *Journal of Research in Science Teaching: The Official Journal of the National Association for Research in Science Teaching*, 34, 377-398, 1997, M. J. Sanger, T. J. Greenbowe.
42. An analysis of college chemistry textbooks as sources of misconceptions and errors in electrochemistry, *Journal of chemical Education*, 76, 853, 1999, M. J. Sanger, T. J. Greenbowe.
43. Misconceptions in Electrochemistry: How Do Pedagogical Agents Help?, *Overcoming Students' Misconceptions in Science: Strategies and Perspectives from Malaysia*, M. Karpudewan, A. N. Md Zain, and A. L. Chandrasegaran Eds. Singapore: Springer Singapore, 91-110, 2017, L. T. Tien, K. Osman.
44. Exploring students' understanding of electrochemical cells using an enhanced two-tier diagnostic instrument, *Research in Science & Technological Education*, 32, 229-250, 2014, A. S. L. Loh, R. Subramaniam, K. C. D. Tan.
45. Students' alternative conceptions in chemistry: A review of research and implications for teaching and learning, 1995, 1995, P. J. Garnett, P. J. Garnett, M. W. Hackling.
46. Can Freshman Science Student Teachers' Alternative Conceptions of Electrochemical Cells' Be Fully Diminished?, *Asian Journal of Chemistry*, 24, 485, 2012, F. Karsli.
47. Inquiry-based laboratory activities in electrochemistry: High school students' achievements and attitudes, *Research in Science Education*, 43, 413-435, 2013, B. Acar Sesen, L. Tarhan.
48. Student misinterpretations and misconceptions based on their explanations of two computer animations of varying complexity depicting the same oxidation-reduction reaction, *Chemistry Education Research and Practice*, 13, 471-483, 2012, D. P. Rosenthal, M. J. Sanger.
49. Using a teaching model to correct known misconceptions in electrochemistry, *Journal of chemical education*, 77, 104, 2000, F. Rogers, P. A. Huddle, M. D. White.
50. The origins of the concept mapping tool and the continuing evolution of the tool, *Information visualization*, 5, 175-184, 2006, J. D. Novak, A. J. Cañas.
51. High school and preservice chemistry teacher education students' understanding of voltaic and electrolytic cell concepts: evidence of Consistent learning difficulties across years, *International Journal of Science and Mathematics Education*, 20, 1859-1882, 2022, S. Rahayu, D. F. Treagust, A. Chandrasegaran.
52. Facilitating conceptual change in students' understanding of electrochemistry, *International Journal of Science Education*, 24, 425-439, 2002, M. Niaz.
53. Factors that prevent learning in electrochemistry, *Journal of Research in Science Teaching: the official journal of the national association for research in science teaching*, 44, 258-283, 2007, H. J. Schmidt, A. Marohn, A. G. Harrison.
54. IMPACT OF INTERACTIVE MULTIMEDIA MODULE WITH PEDAGOGICAL AGENTS ON STUDENTS' UNDERSTANDING AND MOTIVATION IN THE LEARNING OF ELECTROCHEMISTRY, *International Journal of Science and Mathematics Education*, 12, 395-421, 2014, K. Osman, T. T. Lee.
55. Measuring knowledge integration: Validation of four-year assessments, *Journal of Research in Science Teaching*, 48, 1079-1107, 2011, O. L. Liu, H. S. Lee, M. C. Linn.
56. The knowledge integration perspective on learning and instruction, *na*, 2006, M. C. Linn.

57. Supporting knowledge integration in chemistry with a visualization-enhanced inquiry unit, *Journal of Science Education and Technology*, 23, 37-58, 2014, J. L. Chiu, M. C. Linn.
58. Teaching and assessing knowledge integration in science, *Science*, 313, 1049-1050, 2006, M. C. Linn, H.-S. Lee, R. Tinker, F. Husic, J. L. Chiu.
59. Interrater reliability: the kappa statistic, *Biochemia medica*, 22, 276-282, 2012, M. L. McHugh.
60. Scale development: Theory and applications, Sage publications, 2021, R. F. DeVellis, C. T. Thorpe.
61. Inovation of Biology learning through the development of authentic assessment based on scientific literacy for student of senior high school, *Journal of Physics: Conference Series*, 1899, 012142, 2021, N. Bedduside, A. Hadis, A. A. Jalal, D. Patongai.
62. See Supplemental Material for galvanic cell knowledge integration assessment, statistical analysis of the assessment test, and codebook of galvanic cell knowledge integration assessment.
63. A Study of Rasch, partial credit, and rating scale model parameter recovery in WINSTEPS and jMetrik, *Journal of Applied Measurement*, 13, 248-258, 2012, J. P. Meyer, E. Hailey.
64. Item parameter recovery, standard error estimates, and fit statistics of the WINSTEPS program for the family of Rasch models, *Educational and Psychological Measurement*, 65, 376-404, 2005, W.-C. Wang, C.-T. Chen.
65. Critical values for Yen's Q 3: Identification of local dependence in the Rasch model using residual correlations, *Applied psychological measurement*, 41, 178-194, 2017, K. B. Christensen, G. Makransky, M. Horton.
66. Rasch analysis for instrument development: Why, when, and how?, *CBE—Life Sciences Education*, 15, rm4, 2016, W. J. Boone.
67. Evidence of multidimensionality in the ALSFRS-R Scale: a critical appraisal on its measurement properties using Rasch analysis, *Journal of Neurology, Neurosurgery & Psychiatry*, 84, 1340-1345, 2013, F. Franchignoni, G. Mora, A. Giordano, P. Volanti, A. Chiò.
68. Rasch analysis of the Kentucky inventory of mindfulness skills, *Mindfulness*, 7, 466-478, 2016, O. N. Medvedev, R. J. Siegert, P. Kersten, C. U. Krägeloh.
69. Investigating reliability and validity of student performance assessment in Higher Education using Rasch Model, *Journal of Physics: Conference Series*, 1529, 042088, 2020, T. Z. H. T. Petra, M. J. Ab Aziz.

KEYWORDS

knowledge integration, galvanic cell, interdisciplinary learning, physics chemistry

* Corresponding author email: 476485934@qq.com

Research on the Application of Virtual Simulation Technology in Physics Education

¹Bingqi Xue

¹School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (Ministry of Education), Southeast University, Nanjing, Jiangsu 211189, China

ABSTRACT

This study explores the application effectiveness of virtual simulation technology in inquiry-based physics instruction, using the topic of ideal gases in junior secondary school physics as a vehicle. A virtual simulation tool for visualizing the motion of ideal gas molecules was designed and developed, enabling the simulation of molecular motion under adjustable parameters such as temperature and volume. Based on this tool, an inquiry-based lesson titled The Invisible “Gas Spirits” was designed for eighth-grade students, integrating elements such as engaging introduction, foundational knowledge building, virtual inquiry, and real-life applications to establish connections between microscopic and macroscopic understanding. Classroom assessment questions were developed, covering topics such as molecular thermal motion and the relationship between gas pressure, temperature, and volume. Data collected include students’ hands-on operation data, assessment responses, video recordings, and classroom participation performance. Subsequent analysis will focus on the effectiveness of virtual simulation in enhancing students’ conceptual understanding, students’ inquiry abilities when manipulating different parameters, and the practical value of virtual demonstration technology in addressing instructional challenges related to microscopic physics concepts.

KEYWORDS

virtual simulation, physics teaching, gas molecular motion, physics education

TWO ENERGY EXPRESSIONS REFLECT DIFFERENT SYSTEMS UNDERSTANDING

Zihao Yang¹ and Lei Bao^{1,*}

¹College of Design and Engineering, National University of Singapore, Singapore.

²College of Integrative Studies, Singapore Management University, Singapore.

³Engineering Cluster, Singapore Institute of Technology, Singapore.

[leave two blank lines between author affiliations and “abstract”]

ABSTRACT

Energy is an essential concept for physics and science education as it links diverse phenomena through conservation, yet students often describe energy changes by saying that energy “disappears” or is “used up”. These expressions are usually treated as variants of the same misconception. In this study, we test a different hypothesis suggesting that they reflect distinct levels of systems understanding. We analyzed 2,310 item-level responses from think-aloud interviews with Grade 6, Grade 8 and university students ($n = 165$) across 14 everyday energy contexts. Using an interaction-based framework, we coded each response independently for its direct answer to the prompt “Where does the energy go?” and for the systems elements coordinated in the accompanying explanation. Response category and systems-understanding level were strongly associated, and the predicted pairings dominated the joint distribution (Cramer's $V = 0.627$; 73.9% observed versus 28.5% expected under independence). Explicit interaction contexts increased scientifically normative responses overall, but this advantage was explained by stronger systems understanding, especially in Grade 8 when such understanding was solidly forming. After accounting for systems understanding, context was no longer clearly associated with response category. The results show that two classic energy expressions are not equivalent misconceptions, but markers of different representational states with direct implications for assessment and instruction.

KEYWORDS

Energy change, Systems understanding, Misconception, Physics education, Assessment

RESEARCH ON PROMOTING DEEP LEARNING IN HIGH SCHOOL PHYSICS BASED ON COMPLEX PROBLEM SITUATIONS

Xianglong Shan

School of Physical Science and Technology, Soochow University, Suzhou, Jiangsu, 215006, China.

ABSTRACT

In physics education, students learning in complex problem situations has been widely recognized as challenging, with instruction often remaining fragmented and superficial when addressing real-world problems, making it difficult to promote deep learning. To address this deficit, this study develops a teaching model for high school physics driven by complex problem situations, grounded in a conceptual framework that integrates deep learning principles with structured problem-solving processes. The proposed model consists of six interconnected components: creating problem situations, deconstructing problem chains, building learning scaffolds, forming solutions, constructing knowledge systems, and applying knowledge transfer. Guided by this model, we implemented a quasi-experimental design with three rounds of intervention, assessment, and optimization, using a combination of quantitative and qualitative measures to assess students' deep learning levels and thinking structures. The assessment results show that the experimental group significantly outperformed the control group in deep learning motivation, strategies, and thinking structures, particularly at the relational and extended abstract levels of the SOLO taxonomy, as well as in scientific inquiry, collaboration, and problem-solving performance. These findings indicate that the proposed teaching model can effectively represent and enhance students' deep learning development, providing a useful framework for physics instruction aimed at fostering integrated knowledge and higher-order thinking.

REFERENCE STYLE

Cognitive psychology and its implications, W H Freeman and Company, 1980, J.R. Anderson.

Competencies for complexity: Problem solving in the 21st century, Assessment and Teaching of 21st Century Skills: Research and Applications, 41–53, 2018, J. Funke, A. Fischer, D.V. Holt.

Complex problem solving, intelligence, and learning ability, *Complex problem solving: The European Perspective*, 177–200, 1995, J.F. Beckmann, J. Guthke.

Complex problem solving: More than reasoning?, *Intelligence*, 40, 1–14, 2012, S. Wüstenberg, S. Greiff, J. Funke.

Complex problem solving: Toward a (computer-simulated) theory, *Complex problem solving: The European Perspective*, 1995, D. Dörner, A. Wearing.

Complex Problem Solving: What It Is and What It Is Not, *Frontiers in Psychology*, 8, 2017, D. Dörner, J. Funke.

Definitions, traditions and a general framework for understanding complex problem solving, *Complex problem solving: The European perspective*, 3–25, 1995, P. Frensch, J. Funke.

Die Entwicklung eines Modells der Gruppeneffektivität und eines Analyse-Instruments zur Erfassung des Leistungspotentials von Arbeitsgruppen [The development of a group effectiveness model and an analysis instrument for measuring the output potential of work groups], *Verlag Empirische Padagogik*, 2002, P. Simon.

Do complex problem solving dimensions measure something over and above working memory capacity?, *Learning and Individual Differences*, 24, 42–52, 2013, F. Schweizer, S. Wüstenberg, S. Greiff.

Dynamic decision making: Human control of complex systems, *Acta Psychologica*, 81, 211–241, 1992, B. Brehmer.

How does emotion influence different creative performances? The mediating role of cognitive flexibility, *Cognition and Emotion*, 28, 834–844, 2014, W.-L. Lin, P.-H. Tsai, H.-Y. Lin, H.-C. Chen.

Impact of cognitive abilities and prior knowledge on complex problem solving performance - empirical results and a plea for ecological - valid microworlds, *Frontiers in Psychology*, 9, 2018, H.-M. Süß, A. Kretzschmar.

Individual differences in students' complex problem solving skills: How they evolve and what they imply, *Journal of Educational Psychology*, 108, 1028–1044, 2016, S. Wüstenberg, S. Greiff, M.P. Vainikainen, K. Murphy.

Lohhausen. *Vom Umgang mit Unbestimmtheit und Komplexität*, Huber, 1983, D. Dörner, H.W. Kreuzig, F. Reither, T. Stäudel.

On the relationship between task performance and associated verbalizable knowledge, *Q J Exp Psychol*, 36, 209–231, 1984, D.C. Berry, D.E. Broadbent.

Problemlösen in Gruppen: Auswirkungen von psychologischen Trainingsmaßnahmen und Expertenbeteiligung unter kooperativen und kompetitiven Arbeitsbedingungen, *Technische Universität Braunschweig*, 1995, J. Wetzel.

The Logic of Failure, Metropolitan Books, 1996, D. Dörner.

The logic of failure: recognizing and avoiding error in complex situations, *Metropolitan Books*, 1996, D. Dörner.

The multiple faces of complex problems: A model of problem solving competency and its implications for training and assessment, *Journal of Dynamic Decision Making*, 1, 1–14, 2015, A. Fischer, J.C. Neubert.

The role of emotions in complex problem-solving, *Cognition and Emotion*, 19, 1252–1261, 2005, M. Spering, D. Wagener, J. Funke.

Understanding the link between need for cognition and complex problem solving, Contemporary Educational Psychology, 55, 53–62, 2018, J. Rudolph, S. Greiff, A. Strobel, F. Preckel.

KEYWORDS

Physics education, Teaching model, Deep learning, Complex problems, Situational teaching

Achievement, Community, Transformation:

How Travel Grants Impact Physics Careers in the Global South

A.K. Hodari^{1,*}, P. Shastri², J. Beamon-Kiene³

¹STEMScholar, Washington, DC, USA.

²The International Center for Radio Astronomy Research, University of Western Australia, Australia.

³Wilson, NC, USA.

ABSTRACT

The International Union of Pure and Applied Physics (IUPAP) Working Group 5: Women in Physics (WG5) was established as a resolution at the 1999 General Assembly in Atlanta, Georgia, “to survey the present situation and report to the Council and the Liaison Committees, and to suggest means to improve the situation for women in physics”. Since March 2002, WG5 has convened eight triennial events, the International Conference for Women in Physics (ICWIP), and supported yearly travel grants to young women physicists from the Global South.

For this paper we present preliminary data on the impact of IUPAP’s investments in 860 travel grants, with recipient selections made by WG5. Initially, applicants from Eastern Europe dominated the applicant pool; but during the last fifteen years women from the Global South—Argentina, Brazil, India, Kenya, Pakistan—predominated. Herein we will present survey and interview data on the impact of travel grants on recipients’ career trajectories, contextualized within national, regional, and global efforts to increase gender equity in physics.

Survey data are analyzed using basic descriptive statistics, and where possible contrasts were made between subpopulations. Interview data are analyzed using an open coding process, allowing our interpretations to rely on respondents own words. We apply [Angela Johnson](#)’s 2020 model of intersectionality as methodology to center grant recipients as knowers rather than mere sources of data, giving them meaningful roles in naming what matters, in co-constructing interpretation, and in shaping how findings are interpreted. This is a justice-oriented approach that will attend to acts of refusal, resilience, and collective care as important units of analysis.

Findings reveal that travel grants make demonstrative impacts on women’s physics careers. They have engendered community building among women scientists, helped recipients create productive partnerships across national and regional borders, and provided women with leadership opportunities within their home countries.

* Corresponding author email: apriel@stemscholar.co

KEYWORDS

Women, Global South, Physics, Careers, Travel Grants

The Dynamic Process of Human-AI Collaboration among College Students in Solving Open-ended Mechanics Problems

Dechao Lu^{1,2}, Jingbin Xue^{1,2}, Yuyang Qi^{1,2}, Lei Bao^{3†} and Wangyi Xu^{1*}

¹Chinese Education Modernization Research Institute of Hangzhou Normal University (Zhejiang Provincial Key Think Tank), Hangzhou, Zhejiang, China

²School of physics, Hangzhou Normal University, Hangzhou, Zhejiang, China

³School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing, China

ABSTRACT

Recent advances in generative artificial intelligence (AI) have raised important questions about its role in supporting students' problem solving in physics. While much existing research has focused on AI's accuracy in producing solutions, less is known about how students may strategically collaborate with AI tools during authentic problem-solving processes. This study examines undergraduate students' interactions with DeepSeek, a reasoning-oriented large language model, as they engaged in solving physics problems designed with rich context and open-ended tasks. The results reveal clear group differences: low-performing students often relied on AI for direct answers, medium-performing students partially integrated AI feedback, while high-performing students engaged in iterative cycles of questioning, interpretation, and self-construction. These findings suggest that the effectiveness of human-AI collaboration in physics problem-solving depends not only on the availability of AI but also on students' ability to critically and productively engage with AI outputs. The study contributes to a deeper understanding of the dynamics of human-AI problem-solving and offers implications for instructional design aimed at fostering effective AI use in physics education and beyond.

KEYWORDS

Problem solving, Human-AI collaboration, Cognitive offloading

FACIAL EXPRESSION RECOGNITION FOR PROBING STUDENTS' EMOTIONAL ENGAGEMENT IN SCIENCE LEARNING

Yayun Gong¹, Xiaoyu Tang¹, Lei Bao^{2,*}

¹Guangdong Provincial Key Laboratory of Quantum Engineering and Quantum Materials, National Demonstration Center for Experimental Physics Education, School of Physics, South China Normal University, Guangzhou, China.

²School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (Ministry of Education), Southeast University, Nanjing, China.

ABSTRACT

Student engagement in science classroom is an essential element for delivering effective instruction. However, the popular method for measuring students' emotional learning engagement (ELE) relies on self-reporting, which has been criticized for possible bias and lacking fine-grained time resolution needed to track the effects of short-term learning interactions. Recent research suggests that students' facial expressions may serve as an external representation of their emotions in learning. Accordingly, this study proposes a machine learning method to efficiently measure students' ELE in real classroom. Specifically, a facial expression recognition system based on a multiscale perception network (MP-FERS) was developed by combining the pleasure-displeasure, arousal-nonarousal, and dominance-submissiveness (PAD) emotion models. Data were collected from videos of six physics lessons with 108 students. Meanwhile, students' academic records and self-reported learning engagement were also collected. The results show that students' ELE measured by MP-FERS was a significant predictor of academic achievement and a better indicator of true learning status than self-reported ELE. Furthermore, MP-FERS can provide fine-grained time resolution on tracking the changes in students' ELE in response to different teaching environments such as teacher-centered or student-centered classroom activities. The results of this study demonstrate the validity and utility of MP-FERS in studying students' emotional learning engagement.

REFERENCE STYLE

Facial Expression Recognition for Probing Students' Emotional Engagement in Science Learning, *Journal of Science Education and Technology*, 34, 13–30, 2025, X. Tang, Y. Gong, Y. Xiao.

* Corresponding author email: lei.bao@outlook.com

KEYWORDS

Student engagement, Emotional learning engagement, Facial expression recognition, Science teaching

This study focuses on the research progress of semantic analysis models in physics education, with particular attention to their applications in open-ended response analysis, automated scoring and feedback generation, learning process text analysis, as well as instructional resource and curriculum text analysis. With the development of natural language processing, pretrained language models, and large language models, semantic analysis models have gradually expanded from the automatic classification and scoring of students' answers to the identification of conceptual understanding, types of misconceptions, reasoning processes, and learning support needs. Compared with traditional objective assessment methods, these approaches can reveal students' underlying understanding structures in physics learning in a more fine-grained way, thereby providing support for formative assessment, personalized feedback, and instructional decision-making. At the same time, existing studies still face several challenges, including limited interpretability, unstable scoring reliability, insufficient cross-task generalizability, and the need for better educational applicability. Therefore, this paper aims to systematically review the major application pathways, technological development, educational value, and practical limitations of semantic analysis models in physics education, and to provide a theoretical foundation for the subsequent development of tools for diagnosing conceptual understanding based on open-ended responses.

Promoting Deep Learning in Physics with the Conceptual Framework Model of Knowledge Integration

Lei Bao

School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing, 211189, China

lei.bao@outlook.com

Deep understanding of core scientific ideas and cross-cutting concepts in STEM has become a central goal of contemporary science education standards. However, extensive research shows that traditional instruction often falls short in helping students develop deep conceptual understanding in physics. To support assessment and instruction that promote knowledge integration and deep learning, we have developed a new Conceptual Framework model, which represents students' knowledge structures with particular attention to the core ideas and sub-dimensions of a given concept. A Conceptual Framework is a content-specific model that maps the knowledge structures of novices and experts, helping to identify missing connections between core conceptual ideas and other elements within the knowledge structure. Assessments based on the Conceptual Framework model emphasize determining which connections exist in students' knowledge structures, while instruction informed by the model targets the development of more expert-like connections. This talk will introduce the Conceptual Framework model and review recent studies demonstrating promising outcomes from its application. Examples of instructional and assessment approaches based on the model will also be discussed.

Promoting Deep Learning in Physics with the Conceptual Framework Model of Knowledge Integration

Lei Bao


School of Biological Science and Medical Engineering, Research Center for Learning Science, Key Laboratory of Child Development and Learning Science (MOE), Southeast University, Nanjing, 211189, China

lei.bao@outlook.com

Deep understanding of core scientific ideas and cross-cutting concepts in STEM has become a central goal of contemporary science education standards. However, extensive research shows that traditional instruction often falls short in helping students develop deep conceptual understanding in physics. To support assessment and instruction that promote knowledge integration and deep learning, we have developed a new Conceptual Framework model, which represents students' knowledge structures with particular attention to the core ideas and sub-dimensions of a given concept. A Conceptual Framework is a content-specific model that maps the knowledge structures of novices and experts, helping to identify missing connections between core conceptual ideas and other elements within the knowledge structure. Assessments based on the Conceptual Framework model emphasize determining which connections exist in students' knowledge structures, while instruction informed by the model targets the development of more expert-like connections. This talk will introduce the Conceptual Framework model and review recent studies demonstrating promising outcomes from its application. Examples of instructional and assessment approaches based on the model will also be discussed.

307


ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)



ROOM	TOPIC / TRACK	SESSION CODES
303	ASTRONOMY & ASTROPHYSIS (AS) / BIOPHYSICS (BP)	AS1, AS2, BP6, BP8
304	PHYSICS / SCIENCE EDUCATION (PE)	PE1, PE2, PE3, PE4
307	ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)	AMO3, AMO6
308	ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)	AMO1, AMO2, AMO4, AMO5, AMO7, AMO8
309	BIOPHYSICS (BP)	BP1, BP2, BP3, BP4, BP5, BP7
310	STATISTICAL MECHANICS & NONLINEAR PHYSICS (SM)	SM1, SM2, SM3, SM4, SM5, SM6
311	QUANTUM MATERIALS & INFORMATION SCIENCES (QMIS)	QMIS1, QMIS2, QMIS3, QMIS4, QMIS5, QMIS6
324	CONDENSED MATTER & MATERIALS PHYSICS (CM)	CM1, CM3, CM5, CM7, CM9, CM12
325	CONDENSED MATTER & MATERIALS PHYSICS (CM)	CM2, CM4, CM6, CM8, CM10, CM13
326	OCPA AWARD / CONDENSED MATTER & MATERIALS PHYSICS (CM)	AWARD1, AWARD 2, AWARD 3. AWARD 4, CM11, CM14
327	COMPUTATIONAL & MATHEMATICAL PHYSICS (CP) / PLASMA PHYSICS (PL)	CP1, CP2, CP3 (PL), CP4
328	CHEMICAL PHYSICS (CHE)	CHE1, CHE2, CHE3, CHE4, CHE5
329	ACCELERATOR PHYSICS (AC) / HIGH ENERGY & PARTICLE PHYSICS (HEP)	HEP1, HEP2, HEP3, HEP4, HEP5, HEP6, AC1
330	ACCELERATOR PHYSICS (AC) / NUCLEAR PHYSICS (NP) / PLASMA PHYSICS (PL)	AC2, AC3, AC4, NP1, NP2 (PL)

308

ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)



ROOM	TOPIC / TRACK	SESSION CODES
303	ASTRONOMY & ASTROPHYSIS (AS) / BIOPHYSICS (BP)	AS1, AS2, BP6, BP8
304	PHYSICS / SCIENCE EDUCATION (PE)	PE1, PE2, PE3, PE4
307	ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)	AMO3, AMO6
308	ATOMIC, MOLECULAR & OPTICAL PHYSICS (AMO)	AMO1, AMO2, AMO4, AMO5, AMO7, AMO8
309	BIOPHYSICS (BP)	BP1, BP2, BP3, BP4, BP5, BP7
310	STATISTICAL MECHANICS & NONLINEAR PHYSICS (SM)	SM1, SM2, SM3, SM4, SM5, SM6
311	QUANTUM MATERIALS & INFORMATION SCIENCES (QMIS)	QMIS1, QMIS2, QMIS3, QMIS4, QMIS5, QMIS6
324	CONDENSED MATTER & MATERIALS PHYSICS (CM)	CM1, CM3, CM5, CM7, CM9, CM12
325	CONDENSED MATTER & MATERIALS PHYSICS (CM)	CM2, CM4, CM6, CM8, CM10, CM13
326	OCPA AWARD / CONDENSED MATTER & MATERIALS PHYSICS (CM)	AWARD1, AWARD 2, AWARD 3. AWARD 4, CM11, CM14
327	COMPUTATIONAL & MATHEMATICAL PHYSICS (CP) / PLASMA PHYSICS (PL)	CP1, CP2, CP3 (PL), CP4
328	CHEMICAL PHYSICS (CHE)	CHE1, CHE2, CHE3, CHE4, CHE5
329	ACCELERATOR PHYSICS (AC) / HIGH ENERGY & PARTICLE PHYSICS (HEP)	HEP1, HEP2, HEP3, HEP4, HEP5, HEP6, AC1
330	ACCELERATOR PHYSICS (AC) / NUCLEAR PHYSICS (NP) / PLASMA PHYSICS (PL)	AC2, AC3, AC4, NP1, NP2 (PL)

Efimov Effect in Long-range Quantum Spin Chains

Ning Sun, Lei Feng*, and Pengfei Zhang*

Department of Physics, Fudan University, China.

ABSTRACT

When two non-relativistic particles interact resonantly in three dimensions, an infinite tower of three-body bound states emerges, exhibiting a discrete scale invariance. This universal phenomenon, known as the Efimov effect, has garnered extensive attention across various fields. However, it remains an open question how analogous universal few-body physics can emerge in low-dimensional quantum platforms. In this work, we demonstrate that the Efimov effect also manifests in long-range quantum spin chains. The long-range coupling modifies the low-energy dispersion of magnons, enabling the emergence of continuous scale invariance for two-magnon states at resonance. This invariance is subsequently broken to discrete scale invariance for the three-magnon problem, leading to the celebrated Efimov effect. We further discuss generalizations to arbitrary spatial dimensions, where the traditional Efimov effect serves as a special case. Our results reveal universal physics in dilute quantum gases of magnons that can be experimentally tested in trapped-ion systems.

REFERENCE

Efimov Effect in Long-range Quantum Spin Chains, arXiv:2502.20759.

KEYWORDS

Efimov effect, Few-body physics, Long-range spin systems

* Corresponding author email: leifeng@fudan.edu.cn

* Corresponding author email: pengfeizhang.physics@gmail.com

WIDE-BAND MOIRE SUPERCHIRAL OPTICAL FIELD

Qiang Zhang^{1,*}

¹Institute of Microscale Optoelectronics, Shenzhen University, China.

ABSTRACT

On-chip optical superchiral fields are usually generated using nanostructures. Here we demonstrate superchiral field generation using surface waves in a one-dimensional photonic crystal (PhC). The PhC supports nodal rings featuring a bandwidth-enhanced, all-angle degenerate dispersion for transverse magnetic (TM) and transverse electric (TE) surface modes. This unique degeneracy is harnessed to create simultaneous electric field and magnetic field skyrmion bags, resulting in a moiré superchiral field superlattice on the flat crystal surface. We show that the distribution of the superchirality can be modulated by the moiré angle and orbital angular momentum (OAM) of the excitation scheme and the handedness of chirality is determined by the phase difference between TM and TE waves. This work proposes a simple platform for on-chip chiral sensing using tailored superchiral fields.

REFERENCE STYLE

Observation of strong spin-orbit couplings in plasmonic spin-twistronics topological lattices, *Nature Communications*, 17, 1905, 2026, P. Shi, X. Gou, Q. Zhang, W. Wei, H. Wu, Z. Zhu, Y. Shen, X. Yuan.

Bandwidth-enhanced moiré superchiral optical field in a nodal ring photonic crystal, *Physical Review B*, in press, 2026, Q. Zhang, R. Chen, X. Yuan.

KEYWORDS

Near-field optics, Photonic crystals, Chirality

* Corresponding author email: zhangqiang@szu.edu.cn

Stabilizer Rényi Entropy in SYK-like Models

Pengfei Zhang¹

¹Department of Physics, Fudan University, China.

ABSTRACT

Quantum entanglement and quantum magic are two distinct fundamental resources that enable quantum systems to exhibit complex phenomena beyond the capabilities of classical computer simulations. While quantum entanglement has been extensively used to characterize both equilibrium and dynamical phases, the study of quantum magic, typically quantified by the stabilizer Rényi entropy (SRE), remains less explored. In this talk, I will introduce our recent work on investigating the SRE in SYK-like models by representing the corresponding partition function using auxiliary spin degrees of freedom. We apply this method to three examples: (1) the single-site SYK model, (2) the coupled SYK model, and (3) thermofield double dynamics. Our results reveal various transitions in the SRE that are not visible in conventional observables or Rényi entropies of density matrices.

REFERENCE STYLE

- [1]. Stabilizer Renyi Entropy and Its Transition in the Coupled Sachdev-Ye-Kitaev Model, *PHYSICAL REVIEW LETTERS*, 136, 080201 (2026), P. Zhang, S. Zhou, and N. Sun.
- [2]. Connecting Magic Dynamics in Thermofield Double States to Spectral Form Factors, *arXiv:2601.12787*, N. Sun and P. Zhang.

KEYWORDS

Sachdev-Ye-Kitaev model, Quantum magic, Stabilizer Rényi entropy

Application of AI-based methods to quantum metrology and quantum algorithm optimization

Sunny Xin Wang*

Department of Physics, City University of Hong Kong, Hong Kong SAR, China

ABSTRACT

Recent advances in Artificial Intelligence have profoundly influenced science and technology, in particular quantum technologies. In this talk, we present our recent works illustrating how reinforcement learning (RL) and physics-informed neural networks (PINN) can be applied to quantum metrology and quantum algorithm optimization while improving resource efficiency across the quantum workflow.

First, we proposed RL algorithm which can discover generalizable quantum control strategies for parameter estimation tasks. By learning policies that remain effective across a range of parameter values, it enables adaptive control protocols that avoid repeated recalculation of optimal strategies during iterative estimation procedures [1]. Building on this idea, we introduce a reinforcement-learning-based framework for automated circuit architecture discovery in variational quantum sensing. By treating circuit structure as a trainable object and enabling distributed architecture search under hardware constraints, the method adaptively explores state preparation and measurement designs, achieving improved sensitivity and resource efficiency compared with fixed ansatzes [2].

We then present PINN approaches that enhance quantum system characterization and variational algorithm optimization. For Hamiltonian learning, we propose inverse physics-informed neural networks that incorporate the Schrödinger equation into the training objective, enabling accurate inference of system parameters from limited noisy measurements while maintaining physical consistency [3]. Finally, for noisy variational quantum algorithms, we introduce a physics-informed denoising network that learns the effective optimization dynamics under an error-mitigated cost landscape, significantly reducing the number of required circuit executions while preserving the correct optimization trajectory [4].

These results illustrate how AI-driven approaches can improve efficiency across multiple layers of the quantum computing stack and highlight the growing role of AI as a key enabler for scalable quantum technologies.

* Corresponding author email: asim2026@nus.edu.sg [*give email address of the corresponding author in footnote*]

REFERENCE STYLE

[1] Generalizable control for quantum parameter estimation through reinforcement learning, *npj Quantum Inf.*, 5, 82, 2019, H. Xu, J. Li, L. Liu et al.

[2] Automating Variational Quantum Sensing through Reinforcement-Learned Circuit Structures, in preparation, J. Liu, X. Wang.

[3] Hamiltonian learning via inverse physics-informed neural networks, *Phys. Rev. Research*, 7, 043137, 2025, J. Liu, X. Wang.

[4] Accelerating Noisy Variational Quantum Algorithms with Physics-Informed Denoising Networks, in preparation, J. Liu, X. Wang.

KEYWORDS

reinforcement learning, physics-informed neural networks, quantum metrology, quantum algorithm, quantum control

Scalable and efficient generation of graph states in cavity QED

H. H. Jen

Institute of Atomic and Molecular Sciences, Academia Sinica, Taipei, Taiwan.

ABSTRACT

Scalable graph states constitute a fundamental resource for measurement-based quantum computation and a wide range of entanglement-assisted quantum technologies. Their generation requires controllable quantum platforms and efficient, high-fidelity preparation protocols. Here, we propose a systematic approach to generate scalable graph states in one- and two-dimensional geometries using atom–cavity interfaces. Our method is based on the state-carving technique, which enables selective projection of desired entangled states by removing unwanted state components.

We develop a multiqubit state-carving protocol that, combined with the adiabatic transport of a controlled number of atoms in optical tweezers, allows flexible and scalable generation of graph states with tailored geometries. We further analyze the achievable state fidelity and demonstrate that the preparation probability can be optimized through sequential single-photon probing. A key advance of our work is a simple yet powerful modification of the conventional state-carving scheme, which enables entanglement generation with unit probability in principle.

This efficient and experimentally feasible protocol significantly enhances the practicality of entanglement generation in atom–cavity interfaces, even in regimes of moderate cooperativity. Our results establish atom–nanophotonic platforms as a promising route toward large-scale graph-state generation, with direct implications for modular quantum computing, quantum repeaters, and the realization of arbitrarily structured graph states. These capabilities are essential for scalable one-way quantum computation and future quantum network architectures.

REFERENCE STYLE

- [1] Generating graph states in an atom-nanophotonic interface, *Quantum Sci. Technol.* 9, 025020, 2024, C.-H. Chien, S. Goswami, C.-C. Wu, W.-S. Hiew, Y.-C. Chen, and H. H. Jen.
- [2] Efficient and high-fidelity entanglement in cavity QED without high cooperativity, *Phys. Rev. Lett.* 136, 050802, 2026, S. Goswami, C.-H. Chien, N. Sinclair, B. Grinkemeyer, S. Bennetts, Y.-C. Chen, and H. H. Jen.

KEYWORDS

Scalable graph states; Efficient entanglement generation; cavity QED

Quantum magnonics and quantum nonlinear magnonics

Jian-Qiang You*

School of Physics, Zhejiang University, Hangzhou 310027, China

ABSTRACT

Quantum magnonics has emerged as a new direction of research in both quantum physics and condensed matter physics. This talk introduces the background in this area and the very recent progress in my group. First, I will demonstrate the quantum control of a single magnon, including the deterministic generation of a single-magnon state and its coherent superposition with the vacuum (zero-magnon state). Second, I will show some new results on magnon squeezing in the quantum regime, a novel phenomenon in quantum nonlinear magnonics.

REFERENCES

- [1] Quantum Control of a Single Magnon in a Macroscopic Spin System, *Physical Review Letters*, 130, 193603, 2023, D. Xu, X.K. Gu, H.K. Li, Y.C. Weng, Y.P. Wang, J. Li, H. Wang, S.Y. Zhu, J.Q. You.
- [2] Magnon Squeezing in the Quantum Regime, *Nature Communications*, 17, 2679, 2026, Y.C. Weng, D. Xu, Z. Chen, L.Z. Tan, X.K. Gu, J. Li, H.F. Yu, S.Y. Zhu, X. Hu, F. Nori, J.Q. You.

KEYWORDS

Cavity magnonics, Single-magnon control, Magnon squeezing

* Corresponding author email: jqyou@zju.edu.cn

Universal quantum control by time-dependent Hamiltonian:

Hermitian and non-Hermitian

J. Jing¹

¹ School of Physics, Zhejiang University, China.

ABSTRACT

Conventional manipulations over quantum systems for such as coherent population trapping and unidirectional transfer focus on Hamiltonian engineering while regarding the system's manifold geometry and constraint as secondary causes. Here, we treat them on equal footing in controlling time-dependent quantum systems under either Hermitian or non-Hermitian Hamiltonian. The idea is inspired by the D'Alembert principle of regarding active force, constraint force, and inertial force in an unbiased way. For the Hermitian Hamiltonian, we prove that multiple nonadiabatic paths can be obtained by the von Neumann equation for the projection operators of ancillary bases, that serves as dynamical invariants. For the non-Hermitian Hamiltonian, we prove that the Hamiltonian triangularization in the stationary ancillary picture is sufficient to construct at least one nonadiabatic passage in both bra and ket spaces. The passage ends up with a desired target state that is automatically normalized without artificial normalization. Our works demystify and unify the popular protocols on quantum state engineering, including stimulated Raman adiabatic passage, the Lewis-Riesenfeld invariants, quantum transitionless driving, and holonomic transformation. We establish a universal quantum-control framework with a clear motivation inspired by analytical mechanics.

REFERENCE

Universal perspective on nonadiabatic quantum control, *Physical Review A* 111, 012406, 2025, Z. Jin, J. Jing.

Universal quantum control with dynamical correction, *Physical Review A*, 112, 022427, 2025, Z. Jin, J. Jing.

Universal quantum control by non-Hermitian Hamiltonian, *Physical Review A*, 112, 032605, 2025, Z. Jin, J. Jing.

Universal quantum control over bosonic networks, *Physical Review A* 113, 012426, 2026, Z. Jin, J. Jing.

KEYWORDS

universal quantum control, nonadiabatic passage, non-Hermitian Hamiltonian

SU(3) color-orbit coupling in ultracold strontium

C.S. Madasu^{1,2,3}, C. Mitra^{1,2}, L. Gabardos^{1,2}, K.D. Rathod^{2,3}, T. Zanon-Willette⁴, C. Miniatura^{3,5}, F. Chevy^{6,7}, C.C. Kwong^{1,2,*}, D. Wilkowski^{1,2,3}

¹Nanyang Quantum Hub, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore, Singapore.

²MajuLab, International Joint Research Unit IRL 3654, CNRS, Université Côte d'Azur, Sorbonne Université, National University of Singapore, Nanyang Technological University, Singapore, Singapore.

³Centre for Quantum Technologies, National University of Singapore, Singapore, Singapore.

⁴Sorbonne Université CNRS, MONARIS, UMR 8233, Paris, France.

⁵Université Côte d'Azur, CNRS, INPHYNI, Nice, France.

⁶Laboratoire de Physique de l'École normale supérieure, ENS, Université PSL, CNRS, Sorbonne Université, Université de Paris, Paris, France.

⁷Institut Universitaire de France (IUF), Paris, France.

ABSTRACT

Strontium-87, with its zero electronic spin and a large nuclear spin of $9/2$, possesses a large Zeeman manifold within its magnetic-insensitive 1S_0 ground state. This makes ultracold strontium-87 a suitable platform to explore SU(N) physics. We have previously realized uniform SU(2) synthetic gauge field in a tripod configuration, where spin-orbit coupling was observed. Our study reveals particle dynamics in the non-abelian SU(2) gauge field that is similar to Zitterbewegung effect. We extend our study to the SU(3) case by coupling the internal states in a double-tripod configuration. We experimentally demonstrate the all-state connectivity that is inherent in an SU(3) system. As a central result of our work, we study the generalization of spin-orbit coupling to SU(3) color-orbit coupling, where we find three distinct oscillation frequencies with rich dynamics.

REFERENCE STYLE

Experimental realization of a SU(3) color-orbit coupling in an ultracold gas, *Nature Communications*, 16, 8448, 2025, C.S. Madasu, C. Mitra, L. Gabardos, K.D. Rathod, T. Zanon-Willette, C. Miniatura, F. Chevy, C.C. Kwong and D. Wilkowski.

KEYWORDS

Cold atoms, non-abelian gauge field, tripod system, spin-orbit coupling, SU(3)

* Corresponding author email: changchikwong@ntu.edu.sg

Dissipative Phase Transition in a Generalized Quantum Rabi Model

Mingjian Zhu¹, Han Pu^{1,*}

¹Dept. of Physics and Astronomy, Rice University, Houston, TX 77005, USA.

ABSTRACT

We investigate dissipative phase transitions (DPTs) in a parametrically amplified open quantum Rabi model (QRM) with both single- and two-photon decay. In the classical oscillator limit, four composite phases emerge, arising from the possible normal or superradiant regimes across the upper and lower spin branches. A mean-field analysis reveals an “inverted” regime where superradiance emerges only at sufficiently low spin-boson coupling. This regime features first- and second-order DPTs separated by a tricritical point, with two-photon dissipation preserving the stability of the superradiant phase. We further study the steady-state structure beyond the mean-field level and show that the tricriticality stems from the intrinsic nonlinearity of the QRM, unveiled by the interplay of coherent and dissipative two-photon processes. The universality classes of the DPTs are identified, with the corresponding critical and finite-size scaling exponents derived and a scaling ansatz proposed to describe the critical behavior.

REFERENCE STYLE

Dissipative Phase Transition in a Parametrically Amplified Quantum Rabi Model with Two-photon decay, *arXiv:2603.18188*, 2026, M. Zhu, H. Pu.

KEYWORDS

Rabi model, phase transition, superradiance, open quantum system

* Corresponding author email: hpu@rice.edu

FROM NEAR-TERM QUANTUM SIMULATION TO EARLY FAULT-TOLERANCE

Jin Ming Koh^{1,*}

¹Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA.

ABSTRACT

Recent years have witnessed incredible advancements in quantum simulation and computation across both hardware and algorithm fronts. The field stands at an exciting threshold, as an imminent transition from the noisy intermediate-scale quantum era to early fault tolerance promises new capabilities.

In the first part of this talk, I plan to discuss a line of work exploring digital quantum simulation of condensed-matter systems on superconducting quantum processors. Thematically, these studies demonstrate that existing capabilities have already supported scientifically interesting studies. These works probe symmetry-protected topological phases in one dimension [1], interacting Chern insulators hosting chiral modes in two dimensions [2], and higher-order topology in three or more dimensions [3]. The latter studies employ mappings that trade spatial dimensions for interactions, playing to the strength of many-body simulation on a quantum device. More recent work has extended simulation methods to non-Hermitian systems exhibiting anomalous boundary physics [4].

Beyond Hamiltonian physics, digital quantum processors have also enabled direct studies of quantum dynamics. In particular, I highlight a realization of measurement-induced entanglement phase transitions [5], where volume- and area-law phases emerge in random circuits with mid-circuit measurements.

Yet, it has become universally clear that physical noise rates have become limiting, motivating the urgent need for fault-tolerant quantum computation. In the second part of my talk, I therefore introduce a recent work that establishes a class of quantum error-correcting codes that support all-to-all entangling connectivity at near zero cost and perfect fidelity [6]. On state preparation and digital quantum simulation benchmarks, these codes provide substantial logical error rate advantages against the surface code at comparable footprint. This efficiency improvement in error-corrected quantum computation accelerates the timeline to accessing novel, meaningful science beyond classical reach.

* Corresponding author email: kohjm@outlook.com

REFERENCES

- [1] Stabilizing multiple topological fermions on a quantum computer, *npj Quantum Information*, 8, 16, 2022, J.M. Koh, T. Tai, Y.H. Phee, W.E. Ng, C.H. Lee.
- [2] Simulation of interaction-induced chiral topological dynamics on a digital quantum computer, *Physical Review Letters*, 129, 140502, 2022, J.M. Koh, T. Tai, C.H. Lee.
- [3] Realization of higher-order topological lattices on a quantum computer, *Nature Communications*, 15, 5807, 2024, J.M. Koh, T. Tai, C.H. Lee.
- [4] Interacting non-Hermitian edge and cluster bursts on a digital quantum processor, arXiv:2503.14595, J.M. Koh, W.-T. Xue, T. Tai, D.E. Koh, C.H. Lee.
- [5] Measurement-induced entanglement phase transition on a superconducting quantum processor with mid-circuit readout, *Nature Physics*, 19, 1314, 2023, J.M. Koh, S.-N. Sun, M. Motta, A.J. Minnich.
- [6] Entangling logical qubits without physical operations, arXiv:2601.20927, J.M. Koh, A. Gong, A.C. Diaconu, D.B. Tan, A.A. Geim, M.J. Gullans, N.Y. Yao, M.D. Lukin, S. Majidy.

KEYWORDS

Quantum simulation, condensed-matter physics, quantum computation, quantum error correction, quantum dynamics

NOTE TO ORGANIZERS

Due to constraints in my travel plans, I would like to request avoiding scheduling my invited talk on Monday and Tuesday (June 22 and 23), if at all possible.

Assembling, entangling, and probing NaCs molecule arrays

L. Du^{1,2}, C. Williams^{1,2,3}, C. H. Nunez^{1,2,3}, A. Younes^{1,2,3}, K.-K. Ni^{1,2,3}

¹Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, USA

²Harvard-MIT Center for Ultracold Atoms, Cambridge, Massachusetts 02138, USA

³Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

ABSTRACT

Ultracold polar molecules offer an exciting platform for quantum science due to their strong, tunable long-range dipolar interactions and rich internal structure. When assembled into molecule arrays with full quantum control, they can enable the exploration of highly coherent, low-entropy many-body systems in previously inaccessible regimes.

In the Ni group at Harvard, we have been developing a bottom-up approach to assemble, control, and read out ultracold NaCs molecules in optical tweezer arrays starting from individual atoms. In this talk, I will present how ground-state molecules are synthesized in the motional ground state of optical tweezers [1-4], and how long-lived rotational coherence is achieved by engineering the tweezer polarization [5, 6], enabling controlled entanglement between molecules and the realization of an iSWAP gate [7, 8].

I will also discuss recent progress toward non-destructive readout of molecular rotational states using Rydberg atoms, as well as new approaches for scaling to large, defect-free arrays of molecules leveraging electric-field dipolar shielding, paving the way for ultracold molecule arrays to become a scalable quantum simulation platform.

REFERENCE

- [1] Building one molecule from a reservoir of two atoms, *Science* 360, 900 (2018), Lee R. Liu, Jonathan D. Hood, Yichao Yu, Jessie T. Zhang, Nicholas R. Hutzler, Till Rosenband, Kang-Kuen Ni.
- [2] Molecular Assembly of Ground State Cooled Single Atoms, *Phys. Rev. X* 9, 021039 (2019), Lee R. Liu*, Jonathan D. Hood*, Yichao Yu*, Jessie T. Zhang, Kenneth Wang, Yen-Wei Lin, Till Rosenband, and Kang-Kuen Ni.
- [3] Forming a single molecule by magnetoassociation in an optical tweezer, *Phys. Rev. Lett.* 124, 253401 (2020), Jessie T. Zhang, Yichao Yu, William B. Cairncross, Kenneth Wang, Lewis R. B. Picard, Jonathan D. Hood, Yen-wei Lin, Jeremy M. Hutson, Kang-Kuen Ni.

- [4] Assembly of a rovibrational ground state molecule in an optical tweezer, *Phys. Rev. Lett.* **126**, 123402 (2021), William B. Cairncross*, Jessie T. Zhang*, Lewis R. B. Picard, Yichao Yu, Kenneth Wang, Kang-Kuen Ni.
- [5] Rotational Coherence Times of Polar Molecules in Optical Tweezers, *Phys. Rev. Lett.* **127**, 123202 (2021), Sean Burchesky, Loïc Anderegg, Yicheng Bao, Scarlett S Yu, Eunmi Chae, Wolfgang Ketterle, Kang-Kuen Ni, John M Doyle.
- [6] Extended rotational coherence of polar molecules in an elliptically polarized trap, *Phys. Rev. Lett.* **131**, 183401 (2023), Annie J. Park*. Lewis R.B. Picard*, Gabriel E. Patenotte, Jessie T. Zhang, Till Rosenband, Kang-Kuen Ni.
- [7] Site-selective preparation and multi-state readout of molecules in optical tweezers, *PRX Quantum* **5**, 020344 (2024), Lewis R. B. Picard*, Gabriel E. Patenotte*, Annie J. Park*, Samuel F. Gebretsadkan, Kang-Kuen Ni.
- [8] Entanglement and iSWAP Gate between Molecular Qubits, *Nature* **637**, 821 (2025), Lewis R.B. Picard*, Annie J. Park*, Gabriel E. Patenotte, Samuel Gebretsadkan, David Wellnitz, Ana Maria Rey, Kang-Kuen Ni.

KEYWORDS

Ultracold Molecules, Quantum Simulation, Optical Tweezers, Dipolar Shielding

Continuous field tracking with machine learning and steady state spin squeezing

Junlei Duan^{1,*}, Zhiwei Hu¹, Xingda Lu¹, Liantuan Xiao^{2,3}, Suotang Jia^{2,3}, Klaus Mølmer^{4,*}, Yanhong Xiao^{1,2,3,*}

¹Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai, China.

²State Key Laboratory of Quantum Optics and Quantum Optics Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, China.

³Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, China.

⁴Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark.

ABSTRACT

Steady-state entanglement is an important resource for continuous quantum sensing, yet maintaining such entanglement in spin systems remains challenging, particularly in the presence of environmental decoherence and unknown signal perturbations. In this work, we experimentally implement continuous quantum non-demolition measurements with optical pumping on a hot atomic ensemble containing more than 10^{10} atoms, and realize a steady-state spin-squeezed state that can be maintained for more than one day under laboratory conditions.

The sustained spin-squeezed state is then employed for sensing continuously time-varying magnetic fields. To infer the time-dependent signals from the optical measurement records, we utilize recurrent neural network (Long Short-Term Memory) and demonstrated reconstruction of different types of stochastic magnetic fields. The approach demonstrates reliable performance for tracking signals with both Gaussian and non-Gaussian characteristics. Furthermore, we recently implemented deterministic steady state spin squeezing via measurement-based feedback protocol, which enabled quantum-enhanced real-time field tracking. These results provide a pathway toward quantum-enhanced sensing in realistic environments.

REFERENCE

Concurrent spin squeezing and field tracking with machine learning, *Nature Physics*, 21, 909–915, 2025, J. Duan, Z. Hu, X. Lu, L. Xiao, S. Jia, K. Mølmer, Y. Xiao.

KEYWORDS

Spin squeezing, Continuous sensing, Optical magnetometer, Quantum metrology

Occupation-selective topological pumping from Floquet gauge fields

Wenjie Liu^{1,*}, Ching Hua Lee^{2,†}, Zhoutao Lei^{2,‡}

¹School of General Education, Dalian University of Technology, Panjin, China.

²Department of Physics, National University of Singapore, Singapore.

ABSTRACT

Topological pumping is conventionally governed by single-particle band topology. Here we show that promoting tunneling to a dynamical, occupation-conditioned variable fundamentally reshapes this paradigm, leading to occupation-selective topological pumping. In a periodically driven one-dimensional superlattice with density-dependent hopping, two-body bound states (doublons) acquire Chern numbers distinct from those of single particles and exhibit quantized transport even when the single-particle pump is trivial, including counter-propagating responses. We identify a dynamical-gauge-field mechanism that induces topological phase transitions in the bound-state sector absent from the single-particle spectrum. Furthermore, the gauge field concentrates Berry curvature into sharply localized resonant regions without compromising adiabatic quantization. A Floquet realization with ultracold atoms is proposed to realize such occupation-selective pumping. Our results reveal a mechanism for occupation-selective topological responses that can persist across higher-occupancy bound states.

REFERENCE

- [1]Quantization of particle transport, *Phys. Rev. B*, 27, 6083, 1983, D.J. Thouless.
- [2]Thouless pumping and topology, *Nat. Rev. Phys.*, 5, 87, 2023, R. Citro, M. Aidelsburger.
- [3]Progress on new-type optical Thouless pumping, *Laser & Photonics Rev.*, e01528 2025, Y. Chen, Q. Fu, Z. Xu, F. Ye.
- [4]Floquet engineering of correlated tunneling in the Bose-Hubbard model with ultracold atoms, *Phys. Rev. Lett.*, 116, 205301, 2016, F. Meinert, M.J. Mark, K. Lauber, A.J. Daley, H.-C. Nägerl.
- [5]Interaction-induced non-Hermitian topological phases from a dynamical gauge field, *Phys. Rev. Lett.*, 129, 180401, 2022, N. Faugno, T. Ozawa.
- [6]Occupation-selective topological pumping from Floquet gauge fields, arXiv:2603.23307, 2026, W. Liu, C.H. Lee, Z. Lei.

KEYWORDS

Topological pumping, Dynamical gauge field, Doublon states, Floquet systems, Cold atoms

* Corresponding author email: liuwenjie@dlut.edu.cn; phylch@nus.edu.sg; leizht3@nus.edu.sg.

Propagation-invariant spatiotemporal vortices

Junyi Ye^{1,†}, Zhiyuan Che^{1,*,†}, Jiajun Wang¹, Yi Zhang¹, Bo Wang², Wenzhe Liu^{3,*}, Lei Shi^{1,3,4,5,*}, Jian Zi^{1,3,4,5,*}

¹State Key Laboratory of Surface Physics, Key Laboratory of Micro- and Nano-Photonic Structures (Ministry of Education) and Department of Physics, Fudan University, Shanghai 200433, China.

²Henan Key Laboratory of Quantum Materials and Quantum Energy, School of Quantum Information Future Technology, Henan University, Zhengzhou 450046, China.

³Institute for Nanoelectronic Devices and Quantum Computing, Fudan University, Shanghai 200438, China.

⁴Collaborative Innovation Center of Advanced Microstructures, Nanjing University, Nanjing 210093, China.

⁵Shanghai Research Center for Quantum Sciences, Shanghai 201315, China.

† These authors contributed equally to this work.

ABSTRACT

Diffraction, inherently linked to angular spectrum and affected by spatiotemporal coupling effects, imposes a fundamental limit on wave packets propagation. This limitation is particularly acute for spatiotemporal vortex beams, where diffraction distorts the beam profile and splits topological charges. While spatiotemporal coupling engineering has enabled diffraction-free (or propagation-invariant) wave packet propagation, achieving stable vortex propagation remains a significant challenge due to inherent constraints of regular wave dispersion. Here, we overcome this challenge by tailoring the wave system's dispersion to achieve both propagation-invariant behavior and stable vortex structures. We present the first experimental demonstration of propagation-invariant spatiotemporal vortex beams that maintain their spatiotemporal profiles and topological charges over extended propagation distances. The ability of these wave packets to stably carry diverse topological charges without splitting or deformation provides critical insights into spatiotemporal vortex dynamics and unlocks new possibilities for applications across optics, acoustics, and beyond.

REFERENCE STYLE

Propagation-invariant Spatiotemporal Vortices, *Science Bulletin*, 70, 3758-3763, 2025, J. Ye, Z. Che, J. Wang, Y. Zhang, B. Wang, W. Liu, L. Shi, J. Zi.

* Corresponding author email: chezy@fudan.edu.cn (Z. Che), wzliu@fudan.edu.cn (W. Liu), lshi@fudan.edu.cn (L. Shi), jzi@fudan.edu.cn (J. Zi).

KEYWORDS

Spatiotemporal vortices, Propagation-invariant, Spatiotemporal coupling, Liquid surface waves

Macroscopic Spin GHZ State in Levitated Ferromagnets

Xueqi Ni^{1,2,*}, Zhixing Zou¹, Ping Koy Lam^{2,3}, Tao Wang³, Jiangbin Gong^{1,2}

¹ Department of Physics, National University of Singapore, Singapore.

² Centre for Quantum Technologies, National University of Singapore, Singapore.

³ A*STAR Quantum Innovation Centre (Q.INC), Agency for Science, Technology and Research (A*STAR), Singapore.

ABSTRACT

Macroscopic quantum states bridge quantum and classical physics and are central to quantum-enhanced sensing. Among these states, a many-body entangled Greenberger-Horne-Zeilinger (GHZ) state provides maximal entanglement and leads to Heisenberg scaling in quantum sensing. Generating a macroscopic spin GHZ state in a spin system is challenging due to the requirement for coherent control over a large number of particles and complex control sequences. Here, we propose a promising method for generating a macroscopic spin GHZ-like state in a levitated ferromagnet, where the magnetic anisotropy of ferromagnets causes the collective spin and lattice to form strongly dependent degrees of freedom. This strong spin-lattice locking enables mechanical control of the collective spin and defines a constrained Hilbert space where quantum coherence is shared across these coupled degrees of freedom. The resulting GHZ-like state of the collective spins is verified through the Wigner quasi-probability distribution. We further quantify the metrological advantage of the state by demonstrating Heisenberg scaling of the quantum Fisher information. Furthermore, we analyze the roles of symmetry and geometry in decoherence due to gas collisions, identifying experimentally accessible conditions for state realization. Our work addresses the challenge of efficiently generating macroscopic spin GHZ states, reveals a novel form of macroscopic quantum superposition in systems with coupled degrees of freedom, and motivates the platform to test fundamental physics including collapse models into the spin basis.

REFERENCE

- [1] Microscopic theory of a precessing ferromagnet for ultrasensitive magnetometry, *Phys. Rev. Res.*, vol. 7, no. 4, p. 043120, 2025, X. Ni, Z. Zou, R. Lecomte, A. Vinante, D. Budker, P.K. Lam, T. Wang, J. Gong.
- [2] Precessing Ferromagnetic Needle Magnetometer, *Phys. Rev. Lett.*, vol. 116, no. 19, p. 190801, 2016, D. F. Jackson Kimball, A. O. Sushkov, and D. Budker.

KEYWORDS

Macroscopic Superposition, GHZ States, Levitated Ferromagnet, Quantum Sensing

* Corresponding author email: xueqi.ni@u.nus.edu

Large-scale array of squeezed light and synchronization using atomic vapor

L. Wang¹, X. Zhang^{2,3}, K. Manannikov⁴, N. Davidson⁴, Y. Hu^{5,6}, D. Hao^{1*}, Y. Xiao^{1,5,6*}

¹State Key Laboratory of Surface Physics, Key Laboratory of Micro- and Nano-Photonic Structures (Ministry of Education) and Department of Physics, Fudan University, Shanghai, China.

²National Key Laboratory of Infrared Detection Technologies, Shanghai, China.

³Shanghai Institute of Technical Physics, Chinese Academy of Sciences, Shanghai, China.

⁴Department of Physics of Complex Systems, Weizmann Institute of Science, Rehovot, Israel.

⁵State Key Laboratory of Quantum Optics Technologies and Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, China.

⁶Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, China.

ABSTRACT

Quantum light sources such as squeezed light are essential for quantum information science and technologies, but the scalable production of multiple beams of such states remains a challenge. Here, we experimentally demonstrate a novel approach for generating a large spatial array of polarization-squeezed light beams via atomic-coherence-enhanced nonlinear optical processes in a single atomic vapor cell. Unlike schemes based on independent squeezing generators, the squeezing dynamics of each channel are governed here by a common collective ground-state atomic coherence, produced by all input beams, homogenized by the thermal motion of the atoms, and protected against wall collisions by a paraffin coating. Consequently, the optical states of all channels are coupled and mutually regulated via the moving atoms, leading to synchronization behavior. We realized a 30-beam array of polarization-squeezed states with 2.03 ± 0.02 dB of squeezing, experimentally verified the synchronization in both the steady-state and temporal domains, and observed improved purity of the squeezed states as well as the system's response to perturbations, including the temporal response of the squeezing level, as the size of the array increases. This work provides a pathway toward scalable, high-performance quantum light sources for applications in precision measurement, quantum imaging, and quantum information processing.

KEYWORDS

Squeezed state, Optical array, Dissipative coupling

Orientation-Insensitive Spin Amplification for Quantum Metrology

Y. Niu¹, Y. Zhang¹, J. Duan¹, Z. Hu¹, M. Wang², Y. Xiao^{1, 3, 4, *}

¹Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China

²Department of Physics, Wenzhou University, Zhejiang 325035, China

³State Key Laboratory of Quantum Optics Technologies and Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, Shanxi 030006, China

⁴Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, Shanxi 030006, China

ABSTRACT

In quantum metrology based on atom-light interactions, while spin squeezing can be used to reduce atomic projection noise, its metrological gain is often limited by detection noises including photon shot noise. Quantum amplification of the spin signal provides an alternative to spin squeezing, enhancing the system's tolerance to detection noises. However, existing amplification protocols typically restrict the optimal gain to a particular spin direction. Here, we propose a protocol for isotropic quantum amplification of a collective transverse spin and report proof-of-principle experiments in a hot atomic ensemble of $N \sim 10^{11}$ atoms. By combining the vector and tensor atom-light interactions, we engineer an effective Hamiltonian that enables orientation-insensitive spin amplification. By optimizing the amplification process against spin relaxation and decoherences, we experimentally verify its isotropy and obtain about 20% improvement in the signal-to-noise ratio when the system is dominated by detection noises. Our work provides a new perspective for quantum amplification of phase measurements and should find applications in various quantum metrology experiment platforms using collective spins.

KEYWORDS

Quantum Metrology, Quantum Spin Amplification, Orientation-Insensitive

* Corresponding author email: yxiao@fudan.edu.cn

Irreducible Randomness and Apparent Memory in Quantum Stochastic Processes

D. D. C. Chang^{1,2,*}, G. D. Berk^{1,2}, M. Gu^{1,2,3}

¹Nanyang Quantum Hub, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 639673.

²Centre for Quantum Technologies, National University of Singapore, Singapore 117543.

³MajuLab, CNRS-UNS-NUS-NTU International Joint Research Unit, UMI 3654, Singapore 117543.

ABSTRACT

Distinguishing apparent from true irreducible randomness reveals a stochastic process's memory structure—a feature vital for optimal prediction and control. While well-understood classically, quantum analogs remain underdeveloped due to the inherent interactive nature of quantum dynamics. By mapping multi-time input-output relations to temporal Choi states representing quantum stochastic processes, we formulate the quantum entropy rate as a measure of irreducible randomness—expressed as the asymptotic quantum conditional entropy of such temporal states. We prove that causality ensures its non-negativity and convergence of the quantum entropy rate for stationary processes. We show that the quantum entropy rate physically lower bounds the average randomness in the future system evolution no matter how much information an experimenter collects about the process. Furthermore, it directly leads to a quantum excess entropy as the apparent memory which dictates the minimum memory cost for any recurrent quantum circuit model of the process.

KEYWORDS

Quantum Stochastic Processes, Quantum Memory, Non-Markovianity

* Corresponding author email: dere0005@e.ntu.edu.sg

Cold-atom quantum sensing via Bayesian quantum estimation

Chaohong Lee^{1,2*}, Chenying Han^{1,2}, Jiahao Huang³

¹Institute of Quantum Precision Measurement, College of Physics and Optoelectronic Engineering, Shenzhen University, Shenzhen 518060, China.

²Quantum Science Center of Guangdong-Hong Kong-Macao Greater Bay Area (Guangdong), Shenzhen 518045, China.

³Laboratory of Quantum Engineering and Quantum Metrology, School of Physics and Astronomy, Sun Yat-Sen University (Zhuhai Campus), Zhuhai 519082, China.

ABSTRACT

Quantum sensors based on frequentist interferometry face a trade-off between sensitivity and dynamic range. Bayesian quantum estimation, which integrates Bayesian statistics with quantum metrology, overcomes this limitation and enables performance beyond conventional frequentist measurements. In this talk, we present a Bayesian quantum estimation framework for cold atomic ensembles and validate it using cold-atom CPT-Ramsey interferometry. For cold-atom CPT clocks, our adaptive Bayesian protocol achieves Heisenberg-limited sensitivity in integration time and improves fractional frequency stability by 5.1(4) dB over conventional PID locking while enhancing robustness against technical noise [1]. In CPT magnetometry, we optimize measurement sequences to improve precision scaling from $T^{-0.5}$ to $T^{-0.85}$. By tailoring the interferometry sequence via Bayesian quantum estimation, we obtain a dynamic range of 145.6 nT (14.6 dB higher than frequentist counterpart of 5.0 nT) with a sensitivity of 6.8 ± 0.1 pT/Hz^{1/2} (a 3.3 dB improvement over the frequentist counterpart of 14.7 ± 0.4 pT/Hz^{1/2}) [2]. Furthermore, we develop a credible-interval-based adaptive Bayesian quantum frequency estimation protocol for GHZ-state-based atomic clocks, achieving dual Heisenberg-limited precision scaling with respect to both total particle number and total interrogation time [3]. In addition to atomic clocks and magnetometers, this framework may bridge high sensitivity and broad dynamic range for other interferometry-based quantum sensors, such as atomic gravimeters [4].

REFERENCES

- [1] Atomic clock locking with Bayesian quantum parameter estimation: Scheme and experiment, *Physical Review Applied*, 22, 044058, 2024, C. Han, Z. Ma, Y. Qiu, R. Fang, J. Wu, C. Zhan, M. Li, J. Huang, B. Lu, C. Lee
- [2] Adaptive cold-atom magnetometry mitigating the trade-off between sensitivity and dynamic range, *Science Advances*, 11, eadt3938, 2025, Z. Ma, C. Han, Z. Tan, H. He, S. Shi, X. Kang, J., J. Huang, B. Lu, C. Lee

* Corresponding author email: chleecn@szu.edu.cn

[3] Credible-interval-based adaptive Bayesian quantum frequency estimation for entanglement-enhanced atomic clocks, *Science China Physics, Mechanics & Astronomy*, 69, 230312, 2026, J. Zhou, J. Huang, J. Wei, C. Han, C. Lee

[4] Adaptive robust high-precision atomic gravimetry, *Physical Review Research*, 7, L012064, 2025, J. Wei, J. Huang, C. Lee

KEYWORDS

Bayesian quantum estimation, cold-atom quantum sensing, atomic clock, atomic magnetometer, atomic gravimeter

A Robust Laser-Ablated Strontium Cold Atom Source for Compact and Long-Term Operation

Chengmeng Wang^{1,2}, Chang Chi Kwong^{2,3}, David Wilkowski^{2,3,4}

¹Department of Physics, State Key Laboratory of Surface Physics, and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, China.

²School of Physical and Mathematical Science, Nanyang Technological University, Singapore.

³MajuLab, International Joint Research Unit IRL 3654, CNRS, Université Côte d'Azur, Sorbonne Université, National University of Singapore, Nanyang Technological University, Singapore.

⁴Centre for Quantum Technologies, National University of Singapore, Singapore.

ABSTRACT

Cold atom sources are fundamental to the advancement of modern quantum technologies, playing a critical role in fields ranging from quantum computing to precision metrology. However, traditional cold atom platforms, beyond the standard alkaline atoms, typically rely on thermal ovens combined with extensive pre-cooling stages, such as Zeeman slower or two-dimensional magneto-optical traps (2D MOTs). These conventional setups are characterized by large volumes, high power consumption, and demanding maintenance, making them not suitable for widespread deployment in field applications or space-borne quantum sensors where extreme compactness and low energy consumption are required. To address this challenge, we present a compact, laser-induced cold atom source for strontium-88 that eliminates the need for oven, Zeeman slower, and 2D MOT. Our approach utilizes laser ablation, where the focused laser heating a micrometer-sized region compared to the centimeter-sized bulk strontium granule in oven. The bulk material acts as an effective thermal reservoir; once the ablation laser is turned off, the spot cools rapidly, leading to a fast drop in background vapor pressure and the swift recovery of the ultra-high vacuum (UHV) environment. To ensure long-term usability, we implemented an internal mirror to reflect the ablation beam onto the target. This critical design breaks the direct line-of-sight between the strontium vapor and the optical viewports, significantly reducing the probability of atomic deposition and window coating. Furthermore, the mirror prevents the hot atomic flux from directly colliding with the 3D MOT, thereby preserving the trap lifetime. Currently, our highly portable apparatus is capable of loading 1.2 million atoms within 400 ms, achieving a trapping lifetime of 1.5 s. Ongoing optimizations are expected to further enhance the

system's performance, paving the way for next-generation deployable quantum devices.

REFERENCE

Laser-induced thermal source for cold atoms, *Scientific Reports*, 12, 868, 2022, Chung Chuan Hsu, Rémy Larue, Chang Chi Kwong, David Wilkowski.

KEYWORDS

Laser Ablation, Cold Atom Source, Strontium, Magneto-Optical Trap, Compact Setup.

FLOQUET DYNAMICS IN DISPERSIVE AND DISSIPATIVE TIME-VARYING MEDIA

Yuchen Sun¹, Guangwei Hu^{1,*}

¹School of Electrical and Electronic Engineering, Nanyang Technological University, Singapore 639798, Singapore

ABSTRACT

Photonics in media with time-varying properties, such as permittivity, have emerged as promising platforms to break the intrinsic limitations of conventional static photonic systems. The periodic dynamic modulation of the permittivity has unlocked the temporal degree of freedom in the photonic system, thereby giving rise to the generation of Floquet harmonics with frequencies spaced equidistantly by modulation frequency in the spectrum. The periodic dynamic modulation has also facilitated energy exchange between the electromagnetic field and the media, leading to the parametric amplification or coherent absorption of the electromagnetic field. However, the lack of computational techniques has hindered the experimental realization and implementation of the photonic systems with time-varying media.

We start from the first principles, and present a systematic framework for photonic systems with time-varying media. It not only enables previously prohibitive computations, but fosters an insight into the underlying mechanism that provides guidance for the design of realistic photonic systems.

We first focus on the closed Floquet photonic systems, such as photonic time crystal, characterized by the continuous energy bands. The proposed framework enables the fast and robust computation of energy band in complex spatial distributions. This framework further helps reveal novel topological phenomena in the Floquet photonic systems.

In parallel, we focus on the open Floquet photonic scattering systems, such as the leaky cavity, which is essential for the experimental observables. The proposed analytical framework enables modal analysis of the scattering field and establishes the correspondence between the Floquet quasienergy levels and the resonance peaks in spectra. We further introduce a hybrid numerical-analytical simulation algorithm based on the theoretical framework. The open-source software package by the hybrid framework enables the computationally inaccessible simulation of complex Floquet photonic systems with high precision.

REFERENCE

* Corresponding author email: guangwei.hu@ntu.edu.sg

Formulation of dispersive and dissipative time-varying media as a Floquet matrix eigenproblem. *Physical Review Letters* 135, 156903, 2025, Yuchen Sun, Shanhui Fan, and Guangwei Hu.

Quasinormal mode basis for open Floquet photonic systems. *Physical Review Letters*, 2026, Yuchen Sun, Shanhui Fan, and Guangwei Hu. (in press, DOI: 10.1103/175n-lksd)

KEYWORDS

Time-varying photonics, Quasinormal mode, Floquet matrix, Green's function, Scattering theory

Quantum Process Tensors, Combs and Link Product on Infinite Dimensions

A. Lai^{1,*}, G. D. Berk¹, M. Song², M. Gu^{1,2}

¹Nanyang Quantum Hub, School of Physical and Mathematical Sciences, Nanyang Technological University, 637371, Singapore.

²Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, 117543, Singapore

ABSTRACT

Continuous variable (CV) quantum systems can exhibit non-Markovian behavior, and in such cases, a Markovian description discards valuable information that may otherwise have been put to use in the enhancement of quantum control. To alleviate these issues, we extend the framework of quantum combs—a formalism for the characterization and utilization of non-Markovian multi-time processes developed in discrete variable systems—into CV non-Markovian processes. Our key contributions are generalizing the notion of a link product to CV systems and presenting an efficient algorithm to evaluate this for Gaussian processes using covariance matrices. The latter scales polynomially with the number of timesteps of a non-Markovian process, providing a unique advantage over non-Gaussian as well as general DV processes. In doing so, we provide a framework to characterize these non-Markovian processes, and powerful tools with which to study them.

KEYWORDS

Quantum information, quantum foundations, non-Markovian multi-time processes, quantum continuous variables, quantum process tensors

* Corresponding author email: nura0089@e.ntu.edu.sg

Hybrid Optical Lattice-Tweezer Platform for Ultracold Ytterbium Atoms

Bo Song

School of Physics, Peking University, China.

ABSTRACT

Optical lattices and optical tweezers have enabled a series of breakthroughs in quantum simulation and computing. However, conventional optical lattices provide excellent scalability but limited local control, whereas optical tweezers allow flexible and precise manipulation of individual atoms but are limited in system size by the finite field of view of the objective lens.

In this talk, I will present a hybrid optical lattice-tweezer platform for ultracold ytterbium atoms that combines the large-scale scalability of optical lattices with the precise control offered by optical tweezers. This approach enables the preparation and manipulation of large, reconfigurable atomic arrays with high flexibility. I will also report on a two-color magneto-optical trap for ytterbium atoms and rapid optical transport to quantum degeneracy using a moving optical lattice.

Finally, I will discuss our recent progress in fast quantum gas microscopy. In contrast to conventional fluorescence imaging techniques, which typically require exposure times on the order of tens of milliseconds, our approach achieves site-resolved detection with microsecond-scale exposure times. Beyond conventional parity-projection measurements that infer multiple occupancy indirectly, our method enables direct detection of multiple atoms in a single lattice site, paving the way for studying quantum many-body dynamics with high temporal and spatial resolution.

REFERENCE

- [1] Two-color magneto-optical trapping of ytterbium atoms, *Phys. Rev. Applied*, 24, 044039, 2025, X. Li, Y. Wang, L. Yu, B. Song.
- [2] Optical transport of cold atoms to quantum degeneracy, *arXiv:2602.14249*, Y. Tao, Y. Wang, L. Yu, B. Song.

KEYWORDS

Optical Lattice, Optical Tweezer, Ultracold Ytterbium Atoms, Quantum Gas Microscopy.

Long-range nonstabilizerness and quantum codes, phases, and complexity

Fuchuan Wei¹, Zi-Wen Liu^{1,*}

¹Yau Mathematical Sciences Center, Tsinghua University

ABSTRACT

As a necessary resource for quantum computational advantage, quantum magic (nonstabilizerness) is of fundamental importance in the study of quantum computation and physics. We develop a systematic theory of *long-range magic (LRM)*---nonstabilizerness that cannot be erased by shallow unitary circuits---and demonstrate its broad relevance. By bridging LRM with fault-tolerant logical gate theory, we show the emergence of LRM families from quantum error-correcting codes and devise a simple yet powerful method for testing transversal logical gates. Further, we introduce and characterize *LRM phases* in which all ground states throughout the phase exhibit LRM, and identify certain non-Abelian topological orders as representative examples. Then, adopting a complexity theory perspective, we demonstrate the classicality of non-LRM systems in e.g. preparation and learning settings, and present a “no low-energy trivial magic” (NLTM) conjecture with key motivation in the quantum PCP context, for which our LRM results suggest a compelling route. Additionally, we demonstrate how to diagnose LRM with correlation functions. The concepts and results admit substantive extensions to approximate (robust) and nongeometric scenarios. Our LRM theory illuminates profound new connections among quantum resources, computational advantage, error correction and fault tolerance, many-body physics, and complexity theory.

REFERENCE STYLE

Long-range nonstabilizerness and quantum codes, phases, and complexity, arXiv: 2503.04566, F. Wei, Z.-W. Liu.

KEYWORDS

Topological order, Quantum computational resources, Quantum error correction, Quantum complexity theory, Many-body quantum computation

* Corresponding author email: zwliu0@tsinghua.edu.cn

Room-Temperature Vector Atomic Magnetometer and brain magnetic field sensing

J. Lu¹, G. Zhang², X. Meng¹, J. Duan¹, D. Hao¹, K. Zhao^{3,*} and Y. Xiao^{1,4,5,*}

¹Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai, China.

²School of Physics, Zhejiang University of Technology, Hangzhou, China.

³Applied Ion Beam Physics Laboratory, Key Laboratory of the Ministry of Education, and Institute of Modern Physics, Fudan University, Shanghai, China.

⁴State Key Laboratory of Quantum Optics Technologies and Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, China.

⁵Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan, China.

ABSTRACT

Precision measurement of magnetic fields has important applications in brain science, biomedicine, geomagnetic navigation, magnetic anomaly detection etc. Atomic magnetometers are currently the most sensitive tools for magnetic field measurement; however, most high-sensitivity atomic magnetometers are scalar ones that can only measure the magnitude of the field. Vector magnetometers are capable of measuring both the magnitude and the direction of magnetic fields, and therefore can provide more information about the orientation and location of the field source. Yet, due to constraints from the operating principles, most vector magnetometers cannot simultaneously meet the requirements of high sensitivity, real-time, high spatial resolution, single-laser-beam and room temperature for device applications. We demonstrate a room-temperature all-optical vector magnetometer and triaxial brain field sensing using atomic magnetometer based on Electromagnetically induced transparency (EIT) in rubidium vapor cell with anti-relaxation wall coating. Employing the neural-network-based decoding techniques, we achieve the magnetic field amplitude sensitivity of $16.33 \text{ fT}/\sqrt{\text{Hz}}$ and directional sensitivities of $0.015 \text{ mrad}/\sqrt{\text{Hz}}$ for polar angle and $0.032 \text{ mrad}/\sqrt{\text{Hz}}$ for azimuth angle for a magnetic field about 140 nT. We further employ the magnetometer to develop a triaxial magnetoencephalography prototype device, which can sense the brain magnetic field signal in three orthogonal directions of evoked responses from auditory stimulation. Our approach of machine learning assisted vector atomic magnetometry can be extended to high-sensitivity detection of vector geomagnetic field and anomaly magnetic field detection. Meanwhile, our prototype device provides an important technical foundation for room-temperature, non-invasive triaxial bio-magnetic-field measurements.

* Corresponding author email: zhaokf@fudan.edu.cn

* Corresponding author email: yxiao@fudan.edu.cn

KEYWORDS

Atomic magnetometer, Vector magnetometry, Magnetoencephalography,
Electromagnetically induced transparency, Anti-relaxation coating

Electric Bias-induced Reversible Configuration of Single and Heteronuclear Dual-Atom Catalysts on 1T'-MoS₂

Jianhua Wu,^{1,#} Zhongxin Chen,^{1,2,#} Ke Yang,^{3,#} Xin Zhou,^{1,4#} Huizhi Li,⁵ Zhiyong Wang,¹ Mengyao Su,³ Rongrong Zhang,¹ Tie Wang,^{6,7} Qikun Hu,¹ Ning Yan,⁶ Cuibo Liu,⁸ Bin Zhang,⁵ Ming Yang,^{3,*} Shibo Xi,^{9,*} Kian Ping Loh^{1,10*}

¹ Department of Chemistry, National University of Singapore, 3 Science Drive 3, Singapore 117543, Singapore

² School of Science and Engineering, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China

³ Department of Applied Physics, The Hong Kong Polytechnic University, Kowloon, Hong Kong, China

⁴ Institute for Functional Intelligent Materials, National University of Singapore, Singapore

⁵ Department of Chemistry, School of Science, Tianjin University, Tianjin 300072, China

⁶ Department of Chemical and Biomolecular Engineering, National University of Singapore, Singapore 117585, Singapore

⁷ Joint School of NUS and TJU, International Campus of Tianjin University, Fuzhou, 350207, China

⁸ Institute of Molecular Plus, Tianjin University, Tianjin 300072, China

⁹ Institute Of Sustainability for Chemicals, Energy And Environment (ISCE2), A*STAR

¹⁰ Centre for Hydrogen Innovations, National University of Singapore, E8, 1 Engineering Drive 3, 117580 Singapore

ABSTRACT

The development of substrates capable of anchoring single-atom catalysts (SACs) while enabling their dynamic reconfiguration into heteronuclear dual-atom catalysts (DACs) holds significant promise for electrochemical synthesis, yet remains underexplored. Here, we show that electrochemical desulfurization of MoS₂ generates vacancy-rich 1T' domains, which support high loadings of Cu (7.9 wt%) and Pt (6.7 wt%) SACs that are well positioned for dynamic sintering to form DACs. Operando X-ray absorption spectroscopy and density functional theory calculations reveal a voltage-driven, reversible transformation between individual Pt/Cu SACs and Cu-Pt DAC configurations during hydrogen evolution reaction (HER) potentials. The electric field-induced Cu-Pt DACs exhibit superior performance in the selective hydrogenation of alkynes compared to their monometallic SAC counterparts. This work underscores vacancy-enriched 1T'-MoS₂ as a versatile platform for high-density SAC deposition,

enabling on-demand structural reconfiguration and paving the way for tailored catalyst design in electrosynthesis.

REFERENCE STYLE

Electric bias-induced reversible configuration of single and heteronuclear dual-atom catalysts on 1T'-MoS₂. *Nat. Nanotechnol.* 20, 1043–1051, 2025, J.H. Wu, Z.X. Chen, K. Yang, X. Zhou.

KEYWORDS

Bimetallic single-atom catalysts, dual atom catalyst, vacancy engineering, MoS₂, dynamic reconfiguration

Quantum trajectory of a spin oscillator, continuous retrodiction beyond the Heisenberg uncertainty relation

P. Li^{1,2}, J. Duan^{3*}, S. Jia^{1,2}, K. Mølmer^{4*}, Y. Xiao^{3,1,2*}

¹State Key Laboratory of Quantum Optics Technologies and Devices, Institute of Laser Spectroscopy, Shanxi University, Taiyuan, Shanxi 030006, China.

²Collaborative Innovation Center of Extreme Optics, Shanxi University, Taiyuan 030006, China.

³Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China.

⁴Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK 2100 Copenhagen, Denmark.

ABSTRACT

The Heisenberg uncertainty relation imposes a fundamental limit on the measurement precision of non-commuting observables, constraining the performance of quantum sensors and measurement devices. Here, we experimentally demonstrate effectively “simultaneous” sub-SQL (standard quantum limit) estimation of two non-commuting observables in a macroscopic spin oscillator by combining predictive and retrodictive information within the past quantum state formalism. Using a rubidium-87 atomic ensemble of about 10^{10} spins, we perform continuous quantum non-demolition measurements of the canonical quadratures x and p . By synthesizing the forward-evolving quantum state and the backward-propagating effect matrix, we achieve conditional variances for both observables that both lie about 1 dB below the Heisenberg uncertainty bound, with the squeezing maintained over two hours. Moreover, the noise reduction extends to arbitrary quadratures in phase space, all with sub-SQL fluctuations. This work establishes a general framework for surpassing fundamental quantum limits in multi-parameter estimation, with direct applications in quantum-enhanced sensing and continuous-variable quantum information.

KEYWORDS

Quantum trajectory, Past quantum state, Spin oscillator, Heisenberg uncertainty relation, Quantum non-demolition measurement

*Corresponding author email: 20110190074@fudan.edu.cn, klaus.molmer@nbi.ku.dk, yxiao@fudan.edu.cn

A Unified Framework for Sequential Quantum Correlation Sharing

C.L. Ren ^{1,*}, Z.N. Cai ¹

¹ Department of Physics, Hunan Normal University, Changsha 410081, China

ABSTRACT

Quantum measurements not only extract information but also reshape quantum states through measurement backaction, thereby determining how quantum resources can be distributed and recycled. In this talk, recent progress on the sequential sharing of quantum correlations will be reviewed from the perspective of generalized quantum measurements and quantum instruments. Particular emphasis will be placed on Bell nonlocality, EPR steering, entanglement, network nonlocality, and contextuality under sequential and weak-measurement scenarios. Recent advances on complementarity-based correlation sharing and unified measurement frameworks beyond conventional POVM descriptions will also be discussed, highlighting how different Kraus structures govern resource dissipation, robustness, and quantum resource recycling in multipartite and sequential quantum information protocols.

REFERENCE STYLE

- [1] A review of quantum correlation sharing: The recycling of quantum correlations triggered by quantum measurements, *Physics Reports* 1098, 1-53, 2025, Z. Cai, C. Ren*, T. Feng, X. Zhou, J. Chen
- [2] Unified Framework for Quantum Resource Recycling via Instrument-Dependent Back-action, *arXiv:2605.03513*, 2026, Z. Cai, C. Ren*
- [3] Entanglement sharing in sequential quantum measurements revealed by complementarity, *Phys. Rev. A*, 113, 012217, 2026, Z. Cai, C. Ren*
- [4] Nonlocality sharing for a three-qubit system via multilateral sequential measurements, *Phys. Rev. A* 105, 052221, 2022, C. Ren*, X. Liu, W. Hou, T. Feng, and X. Zhou
- [5] Network nonlocality sharing via weak measurements in the extended bilocal scenario, *Phys. Rev. A* 105, 042436, 2022, W. Hou, X. Liu, and C. Ren*
- [6] Observation of nonlocality sharing via not-so-weak measurements, *Phys. Rev. A* 102, 032220, 2020, T. Feng, C. Ren*, Y. Tian, M. Luo, H. Shi, J. Chen, and X. Zhou

KEYWORDS

Quantum correlation sharing, Sequential quantum measurements, Measurement backaction

* Corresponding author email: renchangliang@hunnu.edu.cn

NEW FRONTIERS IN SYNTHETIC GAUGE FIELDS: COMPLEX AND HIGHER-RANK TENSORS

Qi Zhou

Department of Physics and Astronomy, Purdue University, West Lafayette, IN, USA

THE STUDY OF SYNTHETIC GAUGE FIELDS HAS DRIVEN MAJOR ADVANCES ACROSS MANY AREAS OF PHYSICS OVER THE PAST TWO DECADES. MOST OF THESE WORKS, HOWEVER, HAVE FOCUSED ON REAL VECTOR GAUGE FIELDS. IN THIS TALK, I WILL DISCUSS HOW THIS FRAMEWORK CAN BE GENERALIZED TO A LARGELY UNEXPLORED SETTING INVOLVING COMPLEX AND HIGHER-RANK TENSOR GAUGE FIELDS. I WILL PRESENT EXPERIMENTALLY FEASIBLE APPROACHES FOR REALIZING THESE FIELDS AND HIGHLIGHT THE QUALITATIVELY NEW PHENOMENA THAT ARISE WHEN THEY COUPLE TO EXOTIC QUANTUM PHASES.

REFERENCE

Excitonic skin effect, *arXiv:2508.20027*, 2025, W. Xu, Q. Zhou

Synthetic tensor gauge fields, *Physical Review Research* 7, 013013, 2025, J. Zhang, W. Xu, Q. Zhou, S. Zhang

Multipolar condensates and multipolar Josephson effects, *Nature Communications* 15, 4786, 2024, W. Xu, Q. Zhou

Curving the space by non-Hermiticity, *Nature communications* 13, 2184 2022, C Lv, R. Zhang, Z. Zhai, Q. Zhou

KEYWORDS

Gauge fields, macroscopic quantum phenomena, non-Hermitian phenomena

Scalable generation of massive Schrödinger cat states via quantum tunnelling

Han Zhang, Yong-Kui Wang, Yi Zheng, Hai-Tao Bai, Bing Yang.

State Key Laboratory of Quantum Functional Materials, Department of Physics and Guangdong Basic Research Center of Excellence for Quantum Science, Southern University of Science and Technology, Shenzhen, China.

ABSTRACT

Massive spatial quantum superposition states are an important resource for quantum-enhanced metrology and for investigating the interplay between quantum mechanics and gravity. However, generating controllable high-mass spatial superpositions beyond single atoms remains a major experimental challenge. In this work, we report the scalable realization of massive Schrödinger cat states through coherent quantum tunnelling of ultracold atomic clusters in optical lattices. By engineering strongly interacting bosonic atoms in a double-well optical lattice, we observe coherent tunnelling dynamics of bound atomic clusters with an effective mass reaching 608 atomic mass units (AMU). The suppression of tunnelling strength associated with increasing particle number is mitigated through full control of the model parameters. Furthermore, we implement an interferometric protocol to verify the generated spatial entanglement and demonstrate quantum-enhanced phase sensitivity using the resulting cat states. Our results establish a versatile platform for creating and manipulating massive entangled matter-wave states, opening new opportunities for quantum simulation, precision measurement, and experimental studies of macroscopic quantum phenomena.

REFERENCE STYLE

Scalable generation of massive Schrödinger cat states via quantum tunnelling, *Nature Physics*, 2026, H. Zhang, Y.-K. Wang, Y. Zheng, H.-T. Bai, B. Yang.

KEYWORDS

Quantum tunnelling, Schrödinger cat states, Optical lattices, Ultracold atoms, Quantum interferometry.

Wideband pulse shaping for matterwave interferometry and atomic spectroscopy

Yuxiang Zhao^{1,2}, Jinggu Wu^{1,2}, Saijun Wu^{1,2 *}

¹Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China

²Shanghai Key Laboratory of Metasurfaces for Light Manipulation, Shanghai 200433, China

ABSTRACT

In atomic physics, controlling atomic states with light is often challenged by intensity inhomogeneity, while probing atoms with light is constrained by phase information loss. By carefully designing optical pulses at suitable time scales, these constraints can be alleviated, enabling new experimental possibilities. In this talk, I will present two examples, both enabled by wideband optical arbitrary waveform generation (OAWG) [1].

First, by shaping nanosecond pulses according to optimal control theory [2], we achieve $\mathcal{F} \approx 99.2\%$ Raman beam-splitting fidelity with $\pm 50\%$ laser intensity-error resilience. We demonstrate high-contrast, $34\hbar k$ recoil-transfer Raman interferometry operating simultaneously on three pairs of atomic spin states. This technique should be useful for wideband precision inertial sensing in the presence of magnetic field noise.

Second, we develop a frequency-jump approach to coherent atomic spectroscopy. By sending precisely known, rapidly modulated optical waveforms to atoms, wideband spectral information is extracted from transient transmission in a single shot. The method is substantially more data-efficient than traditional frequency-scan methods, ideal for atomic spectroscopy with limited interrogation time window. We demonstrate the technique at an optical nanofiber interface [3], where transmission data is constrained by atomic motion.

REFERENCE

- [1] Superradiant Detection of Microscopic Optical Dipolar Interactions, *Phys. Rev. Lett.* 131, 253602, 2023, Lingjing Ji, Yizun He, Qingnan Cai, Zhening Fang, Yuzhuo Wang, Liyang Qiu, Lei Zhou, Saijun Wu, Stefano Grava, and Darriek E. Chang.
- [2] Composite Biased Rotations for Precise Raman Control of Spinor Matterwaves, *Phys. Rev. Res.* 5, 043094, 2023, Liyang Qiu, Haidong Yuan, Saijun Wu.
- [3] Ferromagnetic Traps for Quasi-Continuous Operation of Optical Nanofiber Interfaces, *Phys. Rev. Applied* 24, 034015, 2025, Ruijuan Liu, Jinggu Wu, Yuan Jiang, Yanting Zhao, Saijun Wu.

KEYWORDS

Atom Interferometry, Robust Quantum Control, Coherent Atomic Spectroscopy

Tailoring Rydberg interactions for expanding quantum capabilities

Q. Liang

Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA

Purdue Quantum Science and Engineering Institute, Purdue University, West Lafayette, Indiana 47907, USA

ABSTRACT

Rydberg atoms provide a uniquely versatile platform for engineering strong, tunable interactions in otherwise weakly interacting neutral-atom systems. They couple strongly to each other and to external fields. The former enables controllable atom-atom, atom-photon and even photon-photon interactions, while the latter promises exceptionally sensitive and broadband microwave (MW) and radio frequency (RF) sensing capabilities. Typically, the Rydberg-Rydberg interactions are leveraged as a blockade mechanism. In contrast, we explore the antiblockade regime in atom arrays, where interactions facilitate rather than suppress excitation. I will present our numerical studies demonstrating directional quantum information and entanglement transfer [1], as well as robust avalanche amplification for weak signal detection [2]. For atom-field coupling, I will present our experimental work expanding MW sensing capabilities with cold atomic ensembles. First, we demonstrate 3D field characterization via multi-parameter extraction, which holds potential as a testbed for machine-learning-enhanced measurement protocols and has broad implications for information extraction from complex measurements. Second, we address the intersection of atom-atom and atom-field interactions by investigating the role of nonlinear atomic interactions in probing external fields [3]. Together, these studies show how tailoring Rydberg interactions expands the accessible measurement and dynamical capabilities of cold-atom platforms.

REFERENCE STYLE

- [1] Yupeng Wang, Junjie Wang, Aishik Panja, Xinghan Wang, Qi-Yu Liang, “Directional Transport in Rydberg Atom Arrays via Kinetic Constraints and Temporal Modulation”, *Phys. Rev. Research* 7, L022035 (2025)
- [2] Xinghan Wang, Yupeng Wang, Qi-Yu Liang, “Robust Rydberg facilitation via rapid adiabatic passage”, *Phys. Rev. Research* 8, 013154 (2026)
- [3] Xinghan Wang, Yupeng Wang, Aishik Panja, Qi-Yu Liang, "Nonlinear optical spectra from Rydberg-mediated photon-photon interactions", *arXiv:2602.11563* (2026)

KEYWORDS

Rydberg atom, electromagnetically induced transparency, quantum information transfer, photon-photon interaction, microwave sensing

Hardware-Aware XZZX Surface-Code Compilation and Dual-Rail Concatenation for Silicon-Atom Architectures

S. Zhang¹

¹Shenzhen international quantum academy, China

ABSTRACT

We study surface-code memory experiments tailored to silicon donor-cluster hardware, where gates are dense within each cluster but sparse and noisier between clusters. Starting from a planar Tanner graph, we compile a rectangular XZZX surface-code patch onto a triangular\rectangular cluster graph with finite cluster capacity, minimizing inter-cluster ancilla-data interactions. Using explicit circuit-level simulations with a silicon-inspired biased noise model, a fixed $d_x = 3$ XZZX family shows decreasing X-memory logical error as d_z increases, with the current 20,000-shot data crossing

an average primitive-noise reference near $d_z \approx 4.69$.

We also extend this framework to simulate dual-rail qubits concatenated with surface codes. The first implementation models each dual-rail block as an effective outer-code qubit with heralded erasure plus residual biased Pauli noise, enabling direct comparison between direct physical-qubit and dual-rail-concatenated surface-code memories. This provides a route to quantify how hardware bias, erasure information, and cluster-aware compilation combine in silicon-compatible fault-tolerant architectures.

REFERENCE STYLE

not published yet

KEYWORDS

QEC, silicon atom qubit, donor-cluster architecture

Most subradiant bound photon pairs from chirality-mediated dispersion softening

Kailin Tan^{1,2,*}, Xuanbing Jiang^{1,2}, Dong Wang^{1,2}, Saijun Wu^{1,2}

¹Department of Physics, State Key Laboratory of Surface Physics and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China

²Shanghai Key Laboratory of Metasurfaces for Light Manipulation, Shanghai 200433, China

ABSTRACT

In the two-excitation sector of waveguide QED, the collective emission from subwavelength atom arrays universally hosts two distinct classes of subradiant eigenstates: free-Fermion states (FSs) and bound photon pairs (bound states, BSs). FSs emerge from the quadratic band extrema of the single-excitation polaritonic dispersion, displaying a universal N^{-3} decay rate scaling with respect to the atom number N ; BSs are subradiant bound atomic excitations that reside within the bandgaps of the scattering continuum. In conventional non-chiral systems, FSs are typically the longest-lived two-excitation states. However, the competition between FSs and BSs in dictating the most subradiant states in chiral waveguide QED, as well as a rigorous analytical derivation for the scaling of BS decay rates, remain largely unexplored.

We address this gap by investigating the subradiant decay rates of BSs in the presence of interaction chirality. We demonstrate that chiral interactions can drive BSs to become the most subradiant two-excitation states across a wide range of lattice spacings. This phenomenon is rooted in a mechanism we term *chirality-mediated dispersion softening*. Specifically, we rigorously prove that the BS decay rate follows the scaling $\Gamma \sim |\alpha_2|/N^3$ if the BS dispersion is $E_K \approx \alpha_2(K-K_{\text{ex}})^2$ near a band extremum K_{ex} , revealing that by increasing chirality, the BS band curvature α_2 can be suppressed, which is the key to achieving extreme subradiance. Aided by an additional toy model, we clarify the general role of chirality for shaping polaritonic dispersion relations toward subradiance, and for engineering exotic photon-photon correlations in nanophotonic systems. Finally, we show that these subradiant chiral BSs can be prepared in a realistic optical nanofiber interface.

REFERENCE

[1] Free-fermion Multiply Excited Eigenstates and Their Experimental Signatures in 1D Arrays of Two-Level Atoms, *Phys. Rev. Lett.*, 128, 093602, 2022, Y.-X. Zhang, K. Mølmer.

* Corresponding author email: kltan22@m.fudan.edu.cn

- [2] Photonic Bound States and Scattering Resonances in Waveguide QED, *arXiv:2110.06093*, 2021, B. Bakkensen, Y.-X. Zhang, J. Bjerlin, A. S. Sørensen.
- [3] Chiral Dissociation of Bound Photon Pairs for a Non-Hermitian Skin Effect, *Phys. Rev. Lett.*, 134, 233602, 2025, J. Shi, A. N. Poddubny.

KEYWORDS

Chiral waveguide QED, Bound photon pair, Subradiant emission, Decay rate scaling

On-chip valley-conserved dual-polarisation photonic crystal router

Kenneth Xin Jie Ho^{1, 2}, Yi Ji Tan^{1, 2}, Ranjan Singh^{3*}

¹Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371

²Centre for Disruptive Photonic Technologies, The Photonics Institute, Nanyang Technological University, Singapore 639798

³Department of Electrical Engineering, University of Notre Dame, Notre Dame, IN 46556, USA

ABSTRACT

Photonic crystals serve as a powerful platform for guiding light in ultra-compact geometries. However, in two-dimensional photonic crystals, overlapping transverse electric (TE) and transverse magnetic (TM) bandgaps cannot be realized with identical permittivities. Here, we achieve overlapping complete photonic bandgaps on a chip by exploiting the difference in 2D effective permittivities for TE and TM modes in lossless silicon photonic crystal slabs with hexagonal air holes arranged in a honeycomb lattice. Leveraging the valley degree of freedom, we design dual-polarization waveguide modes that exist below the light cone and achieve strong field confinement within the slab. This waveguide offers broadband transmission for both polarizations, unlike traditional narrowband TE-only valley photonic crystals with triangular air holes. We further utilize the valley-conserved properties of TE-polarized waveguide modes to design and fabricate an on-chip polarization router through interfacial engineering, employing a mirror-symmetric interface for TE routing and a glide-symmetric interface for exclusive TM propagation. Experimental results demonstrate a 7 GHz region of dual-mode operation within a combined 43 GHz bandwidth (24 GHz TE, 19 GHz TM), together with successful polarization splitting on the chip. Our work presents one of the first experimental realizations of a dual TE/TM polarized valley photonic crystal waveguide and router, enabling polarization-multiplexed functionality for integrated THz photonics. This platform paves the way for high-performance on-chip and inter-chip interconnects, supporting future AI systems, data centre scalability, and emerging 6G communication technologies.

REFERENCE STYLE

On-Chip Valley-Conserved Dual-Polarization Photonic Crystal Router, *APL Engineering Physics*, (in press), 2026, K.X.J. Ho, Y.J. Tan, R. Singh

KEYWORDS

Dual-polarization, valley photonic crystal, THz photonics, on-chip router, polarization multiplexing

Randomized quantum measurements in quantum information processing

H. Zhu

Department of Physics, Fudan University

ABSTRACT

Quantum measurements are the key for extracting information from quantum systems and for connecting the quantum world with the classical world. For many applications in quantum information processing, randomized measurements have proved to be much more efficient in information extraction. In this talk I will discuss the applications of randomized measurements in various tasks, including verification of quantum states, gates, and computation, estimation of linear properties (say fidelity) and nonlinear properties (say purity), certification of (high-dimensional) entanglement, and the learning of unitary channels. In addition, I will explore the efficiency limits of these tasks and highlight the peculiar roles of quantum resources, such as entanglement and nonstabilizerness (magic).

REFERENCE STYLE

1. Qudit Shadow Estimation Based on the Clifford Group and the Power of a Single Magic Gate, Phys. Rev. Lett. 134, 160801, 2025, C. Mao, C. Yi, and H. Zhu.
2. Third moments of qudit Clifford orbits and 3-designs based on magic orbits, arXiv:2410.13575 (accepted by Communications in Mathematical Physics), H. Zhu, C. Mao, and C. Yi.
3. Nonstabilizerness Enhances Thrifty Shadow Estimation, arXiv:2410.23977, D. Chen and H. Zhu.
4. High-Precision Fidelity Estimation with Common Randomized Measurements, Z. Yang, D. Chen, Z. Li, and H. Zhu

KEYWORDS

Randomized quantum measurements, Shadow estimation, Clifford Group, Magic, fidelity estimation

Transient entanglement generation in driven chiral networks beyond the secular approximation

Y. X. Foo¹, K. H. Lim², J.-B. You³, L. C. Kwek^{2,4}, D. Aghamalyan⁵

¹Quantum Science and Engineering Centre, Nanyang Technological University, Singapore.

²Centre for Quantum Technologies, National University of Singapore, Singapore.

³Institute of High Performance Computing, A*STAR (Agency for Science, Technology and Research), Singapore.

⁴National Institute of Education, Nanyang Technological University, Singapore.

⁵SMT Pillar, Singapore University of Technology and Design, Singapore.

ABSTRACT

We study transient entanglement generation between two quantum nodes coupled through a chiral one-dimensional channel. In an emitter-only Born-Markov description, we show that continuous driving and an initial ground state can raise the maximum transient concurrence above the undriven $2/e$ benchmark associated with the effectively single-excitation model [1]. We then consider a more microscopic XX spin-chain channel with triangular plaquette couplings and compare a nonsecular time-convolutionless master equation (TCL-ME) with matrix-product-state (MPS) simulations. In the optimal driven regime, the nonsecular TCL-2 treatment reproduces the concurrence envelope and first transient peak qualitatively, while the remaining discrepancy is mainly attributable to beyond-Born system-bath correlations. The enhancement is traced to the failure of the secular approximation under strong driving, where nearby dressed transitions are not well separated on the dissipative timescale and nonsecular terms mix dressed-state coherences. Finally, we examine within TCL-2 the sensitivity of the protocol to positional disorder, imperfect chirality, and loss into nonguided modes. These results clarify when the familiar $2/e$ limitation ceases to apply and separate the roles of secular breakdown, Born-factorization error, and reduced-state memory in driven chiral entanglement generation.

REFERENCES

1. Chiral route to spontaneous entanglement generation, *Phys. Rev. B*, 92, 155304, 2015, C. Gonzalez-Ballester, A. Gonzalez-Tudela, F. J. Garcia-Vidal, E. Moreno.

KEYWORDS

Waveguide QED, Open Quantum Systems, Entanglement Generation

Microscopic Dynamics in Simulating Lattice Gauge Theory with Ultracold Atoms

Zhen-Sheng Yuan^{1,2,*}

¹University of Science and Technology of China, Hefei 230026, China

²Hefei National Laboratory, University of Science and Technology of China, Hefei 230088, China

ABSTRACT

Exploring the fundamental structure and basic laws of the universe constitutes an essential drive to physicists. The studies of ultracold atoms have built a bridge between the principles of microscopic world and the emergent macroscopic matter world. One can prepare and manipulate synthetic quantum material to simulate strongly correlated quantum many-body system with microscopic techniques to solve formidable tasks for the state-of-the-art supercomputers. I will introduce our recent research on one of such synthetic quantum material, the lattice gauge theory (LGT). We implemented a U(1) LGT Hamiltonian with ultracold atoms trapped in optical lattices and studied the relevant properties of gauge invariance, thermalization dynamics and quantum criticality.

REFERENCES

Paper Title, *Journal Name*, Volume Number, Page Numbers, Year of Publication, Initial1. LastName1, Initial2. LastName2, Initial3. LastName3, Initial4. LastName4.

Example: High-Performance Lithium-Ion Batteries Using Nanostructured Silicon Anodes, *Advanced Energy Materials*, 12, 2103456–2103465, 2022, J.A. Smith, L.B. Chen, M.K. Patel, R.T. Johnson.

- [1] Observation of microscopic confinement dynamics by a tunable topological angle, *Nature Physics* 21, 155 (2025). Wei-Yong Zhang et al.
- [2] Observation of gauge invariance in a 71-site Bose-Hubbard quantum simulator. *Nature* 587, 392 (2020). Bing Yang et al.
- [3] Observation of many-body scarring in a Bose-Hubbard quantum simulator. *Phys. Rev. Res.* 5, 023010 (2023). Guo-Xian Su et al.
- [4] Interrelated thermalization and quantum criticality in a lattice gauge simulator, *Phys. Rev. Lett.* 131, 050401 (2023). Han-Yi Wang et al.
- [5] String-Breaking Mechanism in a Lattice Schwinger Model Simulator *Phys. Rev. Lett.* 135, 101902 (2025). Ying Liu et al.

* yuanzs@ustc.edu.cn

Hilbert Space Fragmentation in a Rydberg Atom Array

Hilbert space fragmentation, where the Hilbert space has been shattered into exponentially many disconnected subspaces, is a form of ergodicity breaking that can lead to rich dynamics in interacting quantum systems. In this talk, I will present our insights on realizing a broad class of Hamiltonians with Hilbert space fragmentation and discuss the implications of the Hilbert space structure on the quantum dynamics of Rydberg atom arrays. First, we uncover generalized quantum many-body scars arising from $\mathbb{Z}_{2k}$ -ordered initial states. In addition, we find that quantum thermalization proceeds in a way that is restricted to a particular Hilbert space fragment. Notably, thermalization between states belonging to different subspaces is precluded, even when these states have the same energy, defying expectations from the eigenstate thermalization hypothesis. Finally, we present the first experimental signatures of statistical localization in a constrained lattice gauge theory model, where expectation values of all conserved quantities remain locally distributed in typical quantum states, even though they are described by nonlocal string-like operators. Our work enables new capabilities on engineering far-from-equilibrium dynamics with control of the Hilbert space structure.

Hilbert space fragmentation, where the Hilbert space has been shattered into exponentially many disconnected subspaces, is a form of ergodicity breaking that can lead to rich dynamics in interacting quantum systems. In this talk, I will present our insights on realizing a broad class of Hamiltonians with Hilbert space fragmentation and discuss the implications of the Hilbert space structure on the quantum dynamics of Rydberg atom arrays. First, we uncover generalized quantum many-body scars arising from Z_{2k} -ordered initial states. In addition, we find that quantum thermalization proceeds in a way that is restricted to a particular Hilbert space fragment. Notably, thermalization between states belonging to different subspaces is precluded, even when these states have the same energy, defying expectations from the eigenstate thermalization hypothesis. Finally, we present the first experimental signatures of statistical localization in a constrained lattice gauge theory model, where expectation values of all conserved quantities remain locally distributed in typical quantum states, even though they are described by nonlocal string-like operators. Our work enables new capabilities on engineering far-from-equilibrium dynamics with control of the Hilbert space structure

A promising platform: from quantum advantage to universal fault-tolerant quantum computing

In recent years, several demonstrations of quantum advantage have drawn tremendous attention to its potential applications in specific areas. However, a critical leap in the quantum technology revolution—from the Noisy Intermediate-Scale Quantum (NISQ) era to universal fault-tolerant quantum computing remains the elephant in the room. This talk will focus on how the strengths of the neutral atom platform can break the “quantity-quality” paradox and our efforts to build a quantum machine with an exceptionally large scale of qubits and high-fidelity quantum gates. Several milestones have been established in our hardware development roadmap, including but not limited to scaling up the number of qubits, optimizing the fidelity of single-qubit and two-qubit gates, improving coherent transport, enabling parallel manipulation, and so on. Meanwhile, in collaboration with our partners, we have designed a blueprint for implementing efficient quantum error correction (QEC) codes (e.g., surface codes, color codes, qLDPC codes, etc.) on our quantum processor, whose layout is closely integrated with GPU-FPGA-QPU architecture.

We experimentally realized optical parametric down-conversion in the Langevin regime on a chip. By precisely controlling the loss inherently tied to fluctuation, we also observed the asymmetric Hong-Ou-Mandel dip—a hallmark of quantum fluctuation in the fluctuation-driven parametric down-conversion—and the fluctuation-induced compression of single photons. Our findings pave the way for manipulating quantum fluctuations, quantum states, and system-reservoir interactions.

Quantum computing platforms—including trapped ions, Rydberg atoms, superconducting circuits, and photonic systems—are rapidly advancing from physical to logical qubits with high-fidelity gates and versatile controllability. In this talk, I present our recent progress in quantum algorithm design for both combinatorial optimization and finite-temperature quantum simulation. In the first part, I address the challenge of encoding NP-hard optimization problems into atomic quantum systems. Using atom-cavity architectures with all-to-all connectivity, we show that the Number Partition Problem, vertex cover, 3-SAT, and general binary optimization (QUBO) can be efficiently mapped onto the cavity-mediated interactions with only linear overhead in atom number—a significant improvement over previous local architectures that require quadratic or cubic scaling. We classify a broad range of NP-complete problems by their encoding complexity into the atom-cavity Hamiltonian, identifying problems with linear, quadratic, and quartic scaling. In the second part, I introduce the Quantum Kernel Function Expansion (QKFE), a method that converts quantum dynamics into quantum thermodynamics. By expanding the density of states in a Fourier series whose coefficients are efficiently measurable on a quantum computer, we obtain the free energy, thermal entropy, and heat capacity as analytic functions of temperature—without variational optimization. I present experimental realizations on the Zuchongzhi superconducting processor using both fully digital and hybrid digital-analogue protocols, demonstrating finite-temperature phase transitions in transverse-field Ising and XY models.

Fault-tolerant resource estimation must be driven by algorithms rather than hardware assumptions alone. In the fault-tolerant regime, different algorithms impose different requirements on logical operations, circuit depth, measurement schedules, magic-state consumption, code distances, runtime, and physical-qubit overhead. This work presents an algorithm-aware view of fault-tolerant resource estimation and organizes the current landscape into several key components: algorithmic workload characterization, logical-level compilation, error-correction overhead, architecture-dependent implementation, and end-to-end cost modeling. We review recent progress in these directions and emphasize that realistic resource estimates require a co-design perspective connecting algorithms, fault-tolerant protocols, and hardware architectures

Trapping over 10,000 individual atoms in a tweezer array by a single metasurface

Wenlan Chen

Tsinghua University

Abstract:

The scalability of physical qubit numbers is a central challenge toward a universal fault-tolerant quantum computer. Here, we robustly trap over 10,000 individual atoms in a tweezer array, enabling the trapped qubit resource to enter the tens-of-thousands scale. This advance is enabled by a single metasurface, 1.98 cm in diameter, that generates the entire tweezer array from a Gaussian beam without the need for microscope objectives, maximizing laser power efficiency. The large aperture of the metasurface ensures a working distance of 1.5 cm, allowing the metasurface to be placed outside the vacuum cell and avoiding the technical complications of in-vacuum operation. We further characterize the randomly loaded atom array using the statistical theory of percolation phase transitions. This work takes an important first step toward a quantum computer at the 10,000-qubit scale.

In the second part, I will show our new setup integrating high-cooperativity cavity with atomic array. Our millimeter-scale Fabry-Pérot cavity reaches the single-atom cooperativity of 125, with high resolution microscopes ($NA=0.65$) for addressing and imaging individual atoms in optical tweezers in the cavity mode. Atom-cavity transmission spectra of trapped atoms yield an effective spectroscopic cooperativity above 110, providing an in-situ verification of strong coupling in the integrated platform with up to 16 individually trapped atoms at the anti-node of the cavity mode. This setup enables us to investigate entanglement with programmable many-body interactions.